

SPECTRAL CALCULATIONS ON LOCALLY CONVEX VECTOR SPACES

GYULA LAKOS

ABSTRACT. We develop a holomorphic functional calculus for (multivalued linear) operators on locally convex vector spaces. This includes the case of fractional powers along Lipschitz curves.

CONTENTS

Introduction	1
1. Lipschitz curves and curve systems	3
2. Integrals of Cauchy type	4
3. Certain estimates and applications to Cauchy integrals	7
4. Rational functions and related identities	14
5. Normal forms of slowly growing holomorphic functions	18
6. Arithmetics of multivalued linear operators	23
7. The holomorphic functional calculus	28
8. Logarithms and fractional powers	29
9. Convergence improvement	32
References	32

INTRODUCTION

If A is a not everywhere defined operator on a locally convex vector space $\mathfrak{V}$ and Γ is a Lipschitz curve system on the Riemann sphere then $u \in D_\Gamma^n(A)$ if $u \in \mathfrak{V}$ and the resolvent operators of A with poles from Γ applied n times to u give a continuous function on Γ^n . The slowly growing holomorphic functions on the complement of the Lipschitz curve system Γ form the algebra $\mathcal{H}_\Gamma$. This algebra has a natural filtration: $f \in \mathcal{H}_\Gamma^n$ if f is a superposition of rational functions with a pole in Γ with multiplicity at most n . In case $\Gamma \subset \mathbb{C}$ it means that

$$f(z) = \sum_{i=0}^n \int_{w \in \Gamma} \frac{\mu_i(w)}{(w - z)^i},$$

where μ_i are finite Borel measures on Γ . Having such representations we may define the operator $f(A)$ on $D_\Gamma^n(A)$ if $f \in \mathcal{H}_\Gamma^n$. We do that by substituting A into z formally. In the special case above

$$f(A)u = \sum_{i=0}^n \int_{w \in \Gamma} \frac{\mu_i(w)}{(w - A)^i} u.$$

2000 *Mathematics Subject Classification*. Primary: 47A60, Secondary: 47A06.

Key words and phrases. Functional calculus, locally convex vector spaces, Lipschitz curves.

The objective of this paper is to elaborate the notions used above and demonstrate the viability of this functional calculus in arithmetical sense. That includes the statements of Well-definedness, Multiplicativity, and Locality, which hold under realistic conditions.

Inspiration for this work came from three sources: one is the author's interest in taking square roots along special curves in locally convex algebras, cf. [6]; another one is the classical theory of fractional powers of operators with radial spectral cuts, see Komatsu [4], or the book of Martínez Carracedo and Sanz Alix [7]; and the third one is the very classical Riesz functional calculus, cf. Dunford and Schwartz [2], VII.3. We will see how these fit into our picture. For a comprehensive review of related matters see the book of Haase [3]. Although this latter one concentrates on the sectorial case but it is exhaustive in that regard.

In terms of the construction of the calculus outlined above, we can notice that the relationship between the function $f(z)$ and the not everywhere defined operator $f(A)$ is not immediate. Although we want $f(A)$ to specialize to $f(z)$ for scalars, we obtain $f(A)$ through certain measures μ_i :

$$\boxed{\begin{array}{c} \text{slowly growing} \\ \text{holomorphic function } f \end{array}} \rightarrow \boxed{\begin{array}{c} \text{normal form} \\ (\text{measures } \mu_i) \end{array}} \rightarrow \boxed{\begin{array}{c} \text{not everywhere defined} \\ \text{operator } f(A) \end{array}}$$

This character also appears in the structure of the paper: Sections 1–5 comprise a function theoretical discussion. In Sections 1 and 2 we go through a short discussion of Lipschitz curve systems and some elementary properties of integrals of Cauchy type. Section 3 is about some finer properties. The main purpose of this section is that we can formulate the statement of locality later in a slightly stronger form than it would be easier otherwise. So despite its relative length one can skip this section without much consequence. Section 4 is a reminder to some elementary identities, which, in fact, will be generalized later, but they are included here for the sake of better readability. Section 5 is about normal forms of slowly growing holomorphic functions outside a Lipschitz curve system Γ . Practically, it says that internal distributions on Γ (whatever they are) are in bijective correspondence to slowly growing holomorphic functions as boundary values, cf. Berenstein and Gay [1], 3.6. The function theoretical part is relatively lengthy but it turns out that once we properly understand what we want to generalize the task itself is surprisingly easy. In Sections 6–9 the functional calculus is actually discussed. In Section 6 we restate the statements of Section 4 in greater generality, this time applied to multivalued linear operators. The key observation is that the resolvent identity remains valid, so we can generalize everything. In Section 7 we define the function calculus and we discuss its basic properties. In Section 8 we examine that how our functional calculus applies to fractional powers. In Section 9 we mention some modifications which can be applied.

In terms of conventions, in what follows “measure” means Borel measure of bounded variation. They are always considered on T_2 , M_2 , locally compact topological spaces unless noted otherwise. We remark that such measures can be well approximated by discrete measures in weak sense, and integration is a question of sequential convergence. In the first part we consider complex measures and scalar integrals. But eventually, in the second part we also consider vector measures or rather vector integrals, depends on viewpoint. One can take a look at e. g. [5] in order to see how to develop a sufficient integration theory along such measures, but the reader is rather assumed to have working knowledge regarding vector integration.

I would like to thank Markus Haase for some useful comments.

1. LIPSCHITZ CURVES AND CURVE SYSTEMS

1.1. Let $\gamma : [a, b] \rightarrow \mathbb{C}$ be a simple continuous curve, $\Gamma = \gamma([a, b])$. We say that γ is a Lipschitz curve if Γ is locally the graph of a Lipschitz function. We say that γ is a short Lipschitz curve if Γ is actually the graph of a Lipschitz function. In the latter case we say that the vector ξ is transversal in angle $\geq \theta$ if for any $w_1, w_2 \in \Gamma$, $w_1 \neq w_2$ the angle between ξ and the line through w_1 and w_2 is $\geq \theta$. If we introduce a coordinate system (u, v) such that its ordinate points into the direction of ξ then we find that the curve has shape $(u, \varphi(u))$ where $\frac{|\varphi(u_1) - \varphi(u_2)|}{|u_1 - u_2|} \leq \cot \theta$.

From the definition it is clear that being a Lipschitz curve is independent of the parametrization. But bi-Lipschitz parametrizations can always be chosen, ie. for which there exist constants $C_1, C_2 > 0$ such that $C_1|t_2 - t_1| \leq |\gamma(t_2) - \gamma(t_1)| \leq C_2|t_2 - t_1|$. For example, the arc-length parametrization is always like that. It is not hard to see that a curve $\gamma : [a, b] \rightarrow \mathbb{C}$ is a Lipschitz curve with a bi-Lipschitz parametrization if its derivative exist in $\mathcal{L}^1$ sense and its representative function γ' can be chosen so that any $t \in [a, b]$ has a neighborhood $\mathcal{V}$ such that the set $\gamma'(\mathcal{V})$ is contained in a bounded, closed convex set not containing 0. In what follows we always consider Lipschitz curves with bi-Lipschitz parametrizations.

If $t_0 \in [a, b]$ then we define the set of curve directions at $\gamma(t_0)$ as $\overline{\lim_{\substack{t \rightarrow t_0 \\ t \in [a, b]}} \frac{\gamma(t) - \gamma(t_0)}{|\gamma(t) - \gamma(t_0)|}}$.

1.2. Let $\Gamma = \gamma([a, b])$. Going counterclockwise, the analytic boundary $\Gamma^\pm$ of $\mathbb{C} \setminus \Gamma$ in $\mathbb{C}$ contains the endpoint $\gamma(b) = \gamma(b)^+ = \gamma(b)^-$, the left boundary $\Gamma^+ = \{\gamma(t)^+ : t \in (a, b)\}$, the endpoint $\gamma(a) = \gamma(a)^+ = \gamma(a)^-$, and the right boundary $\Gamma^- = \{\gamma(t)^- : t \in (a, b)\}$. If U is a subset of Γ then we use the notation $U^+ = \{u^+ : u \in U\}$, and similarly with U^- and $U^\pm$. In particular, if $\mathcal{U} \subset \mathbb{C}$ is open then the analytic boundary $(\Gamma \cap \mathcal{U})^\pm$ of $\mathcal{U} \setminus \Gamma$ in $\mathcal{U}$ will be an open subset of the analytic boundary $\Gamma^\pm$ of $\mathbb{C} \setminus \Gamma$ in $\mathbb{C}$.

If $f : \mathcal{U} \setminus \Gamma \rightarrow \mathbb{C}$ is a continuous function then we can consider its maximal continuous extension $\hat{f}$ to the analytic boundary of $\mathcal{U} \setminus \Gamma$ in $\mathcal{U}$ such that the domain of the extension intersects the analytic boundary in an open subset. In that case we define f^+ and f^- such that $f^+(z) = \hat{f}(z^+)$, and $f^-(z) = \hat{f}(z^-)$, wherever $\hat{f}$ makes sense.

Lemma 1.3. *Suppose that $\mathcal{U}$ is an open set, $f : \mathcal{U} \setminus \Gamma \rightarrow \mathbb{C}$ is holomorphic and it extends to the analytic boundary $(\mathcal{U} \cap \Gamma)^\pm$. Then, we claim,*

$$\bar{\partial}f = \gamma_*\nu$$

in distributional sense on $\mathcal{U}$, where the measure

$$\nu(t) = \frac{f^-(\gamma(t)) - f^+(\gamma(t))}{2i} d\gamma(t).$$

Proof. Slightly prolonging γ , if necessary, we can forget about the endpoints. Furthermore, the question is local, and we can assume that γ is locally of shape $\gamma(t) = t + i\varphi(t)$. Then we can define the functions $f_\varepsilon(x + yi) = f(x + yi - \varepsilon \operatorname{sgn}(y - \varphi(x)))$ locally. For such functions the statement is a consequence of Green's theorem, because the first derivatives extend to the analytic boundary continuously. The general case follows from taking the limit $\varepsilon \searrow 0$. $\square$

1.4. We say that the simple curves $\gamma_i : [a_i, b_i] \rightarrow \mathbb{C}$ ($1 \leq i \leq p$) and the points w_j ($1 \leq j \leq q$) form a curve system if the sets $\gamma_i([a_i, b_i])$ are pairwise disjoint except at endpoints, the w_j are distinct and they do not lie in any of $\gamma_i([a_i, b_i])$. Then $\Gamma =$

$\{w_1, \dots, w_q\} \cup \bigcup_i \gamma_i([a_i, b_i])$ is the supporting set of the curve system. The endpoints of the γ_i and the points w_j are the nodes of the curve system, the elements of the $\gamma_i([a_i, b_i])$ are the regular points.

If the curves γ_i are Lipschitz curves then we say that the curve system is a piecewise Lipschitz curve system. A curve $\gamma : [a, b] \rightarrow \mathbb{C}$ is a piecewise Lipschitz curve if it can be cut into Lipschitz pieces. This happens if and only if $\gamma([a, b])$ is the supporting set of a piecewise Lipschitz curve system.

We say that a curve system is a Lipschitz curve system if it is piecewise Lipschitz and $\gamma_i([a_i, b_i]) \cap \gamma_j([a_j, b_j]) \neq \emptyset$ implies that $\gamma_i([a_i, b_i]) \cup \gamma_j([a_j, b_j])$ is the supporting set of a Lipschitz curve, ie. the components of the curve system join as pieces of Lipschitz curves. If it causes no confusion then we rather call the supporting set Γ of a Lipschitz curve system as the Lipschitz curve system.

From Lemma 1.3 one can immediately deduce

Lemma 1.5. *Suppose that Γ is a Lipschitz domain, $\mathcal{U}$ is an open subset of $\mathbb{C}$, and f is a holomorphic function on $\mathcal{U} \setminus \Gamma$. Suppose that f extends to the analytic boundary of $\mathcal{U} \setminus \Gamma$ in $\mathcal{U}$ continuously. Then we claim,*

$$\bar{\partial}f = \gamma_*\nu$$

in distributional sense on $\mathcal{U}$, where in regular points the measure ν is locally given as

$$\nu(t) = \frac{f^-(\gamma(t)) - f^+(\gamma(t))}{2i} d\gamma(t),$$

such that $\gamma(t)$ is a local parametrization of Γ , and ν extends to the additional, finitely many nodes of Γ as the zero measure (ie. ν is a continuous measure). $\square$

The relationship of Lemma 1.3 and Lemma 1.5 is typical for what follows: If we have a result concerning Lipschitz curves then it extends to Lipschitz curve systems with appropriate modifications.

1.6. Using local charts Lipschitz curves and curve systems can be defined on any $\mathcal{C}^1$ 2-manifold M . If M is oriented then left and right transversal directions, etc. can also be defined. For us only the case $M = \overline{\mathbb{C}}$, the Riemann sphere, will be of importance.

1.7. An open subset $\mathcal{U} \subset \overline{\mathbb{C}}$ is a Lipschitz domain if there is a Lipschitz curve system $\Gamma \subset \overline{\mathbb{C}}$ such that $\mathcal{U}$ is a connected component of $\overline{\mathbb{C}} \setminus \Gamma$. We can consider the analytic boundary $\partial_{\text{an}}\mathcal{U}$ of $\mathcal{U}$ in $\overline{\mathbb{C}}$. The analytic boundary $\partial_{\text{an}}\mathcal{U}$ and the ordinary boundary $\partial\mathcal{U}$ has the same number of components, but the analytic boundary is richer, there is a surjective continuous map $f : \partial_{\text{an}}\mathcal{U} \rightarrow \partial\mathcal{U}$. Topologically, each component of $\partial_{\text{an}}\mathcal{U}$ is either a point or a circle. In fact, for each connected component K_i of $\partial_{\text{an}}\mathcal{U}$ we can find an oriented boundary curve $\hat{\gamma}_i : [a, b]_{a \sim b} \rightarrow K_i$, such that a.) $[a, b]_{a \sim b}$ is a possibly degenerate closed interval $[a, b]$ with the endpoints identified, b.) the underlying curve $\gamma_i = f \circ \hat{\gamma}_i$ is a piecewise Lipschitz curve, although not necessarily simple, c.) $\mathcal{U}$ “lies on the left” of $\hat{\gamma}_i$, in particular in case of a bounded $\mathcal{U}$ it goes counterclockwise. These oriented boundary curves are unique up to bi-Lipschitz reparametrizations.

2. INTEGRALS OF CAUCHY TYPE


2.1. Let $\gamma : [a, b] \rightarrow \mathbb{C}$ be a Lipschitz curve. We define the associated logarithm function as

$$\log(\gamma, z) = \int \frac{d\gamma(t)}{\gamma(t) - z} = \log|\gamma(b) - z| - \log|\gamma(a) - z| + i \int_{t=a}^b d \operatorname{Arg}(\gamma(t) - z).$$

In fact, $\log(\gamma, z) = \widetilde{\log} \frac{\gamma(b) - z}{\gamma(a) - z}$, where $\widetilde{\log}$ is the unique branch of logarithm such that the function vanishes at ∞ . Regarding local behaviour, for $z \sim \gamma(s)$, $s \in (a, b)$ we see that

$$(1) \quad \log(\gamma, z) = \widehat{\log} \frac{\gamma(b) - z}{z - \gamma(a)} + \operatorname{sgn}(\gamma, z) \pi i,$$

where $\widehat{\log}$ is an appropriate branch of $\log$, and $\operatorname{sgn}(\gamma, z)$ is $+1$ or -1 if z on the left or right side of γ , respectively.



Considering (1) it seems reasonable to extend the logarithm function to $z = \gamma(s) \in \gamma((a, b))$ by taking $\operatorname{sgn}(\gamma, z) = 0$. Indeed, we do so but we do not really consider $\gamma(s)$ to be in the domain of $\log(\gamma, z)$ but we use $\log(\gamma, \gamma(s))$ just as a special expression when a formula calls for it. Notice that $\log(\gamma, \gamma(s))$ can be understood as the Cauchy principal value of the defining integral of $\log(\gamma, z)$.

2.2. More generally, consider a finite Borel measure μ on $[a, b]$, and set

$$f(z) = \int \frac{\mu(t)}{\gamma(t) - z}.$$

Then the function $f(z)$ has the following properties:

- a.) $f(z)$ is holomorphic on $\mathbb{C} \setminus \Gamma$, $f(z)$ vanishes at ∞ .
- b.) $|f(z)| \leq \|\mu\| \operatorname{dist}(z, \Gamma)^{-1}$.
- c.) $f(z)$ is a function of class $\mathcal{L}_{\text{loc}}^p$ ($1 \leq p < 2$).
- d.) $f(z)$ can be understood in distributional sense, and then

$$-\pi^{-1} \bar{\partial} f = \gamma_* \mu.$$

In particular, if such functions $f(z)$ are given then the measure μ can be recovered.

- e.) If $s \in [a, b]$ and η stays in a closed sector not containing any curve direction of γ at $\gamma(s)$ then

$$\lim_{\eta \rightarrow 0} \eta f(\gamma(s) + \eta) = \mu(\{s\}).$$

- f.) Suppose that $I = [a, b] \cap (u, v)$. If f continuously extends to the analytic boundary $\gamma(I)^\pm$ then

$$\mu|_I(t) = m(t) d\gamma(t)$$

where

$$(\dagger) \quad m(t) = \frac{f^+(\gamma(t)) - f^-(\gamma(t))}{2\pi i}.$$

In particular, m is continuous.

g.) Assume that $(a_1, b_1) \subset [a, b]$ and $\mu|_{(a_1, b_1)}(t) = m(t) d\gamma(t)$ where m is Hölder-continuous on (a_1, b_1) . Then f extends to $\gamma((a_1, b_1))^\pm$ continuously. In particular, $(\dagger)$ holds on (a_1, b_1) .

Generalizing the case of the logarithm function here it is reasonable to set

$$f(\gamma(t)) = \frac{f^+(\gamma(t)) + f^-(\gamma(t))}{2}.$$

One can see that $f(\gamma(t))$ is the Cauchy principal value of $\int_{t=a}^b \frac{\mu(t)}{\gamma(t) - z}$. Again, this is a reasonable notation, but we do not consider $\gamma(t)$ to be in the domain of $f(z)$.

h.) Assume that $[a, b_1] \subset [a, b]$ and $\mu|_{[a, b_1]}(t) = m(t) d\gamma(t)$ where m is Hölder-continuous on $[a, b_1]$. Then

$$f(z) = m(a) \log(\gamma, z) + \tilde{f}(z),$$

where $\tilde{f}(z)$ extends continuously to $\gamma([a, b_1))^\pm$. Similar statement holds around b .

i.) More generally, if m is a piecewise Hölder continuous function, then the picture is pretty much the same except $f(z)$ may have logarithmic singularities around the discontinuities of m .

Proof. (a) and (b) are immediate from the definition. (c): On $\mathbb{C} \setminus \text{supp } \gamma$ the function $f(z)$ is given as the convolution of the function $z \mapsto -\frac{1}{z}$ and the compactly supported finite measure $\gamma_*\mu$. The function $z \mapsto -\frac{1}{z}$ is of class $\mathcal{L}_{\text{loc}}^p$ ($1 \leq p < 2$), hence we see that $f(z)$ is a function of class $\mathcal{L}_{\text{loc}}^p$ ($1 \leq p < 2$). (d): This convolution can also be understood in distributional sense. We know that the function $z \mapsto \frac{1}{\pi z}$ is a fundamental solution of the $\bar{\partial}$ operator; from which $-\pi^{-1} \bar{\partial} f(z) = \bar{\partial} * (z \mapsto \frac{1}{\pi z}) * \gamma_*\mu = \gamma_*\mu$ follows. (e) follows from a simple estimate. (f) follows from Lemma 1.3 and point (d). (g): We can assume that γ is a short Lipschitz curve with the direction η transversal, and m is Hölder-continuous on $[a, b]$. Then for $t \sim s$, $\tilde{t} \sim 0$, $\tilde{t} \neq 0$ we see that

$$f(\gamma(t) + \eta \tilde{t}) = m(t) \log(\gamma, \gamma(t) + \eta \tilde{t}) + \int_{u=a}^b \frac{m(u) - m(t)}{\gamma(t) + \eta \tilde{t} - \gamma(u)} d\gamma(u).$$

From that the local behaviour is immediate. (h): If $m(a) = 0$ then $f(z)$ extends continuously to the analytic boundary of $\overline{\mathbb{C}} \setminus \Gamma$ for $z \sim \gamma(a)$. One sees this immediately by slightly prolonging γ at a and extending m as identically zero. Otherwise, we can write $m(z) = m(a) + (m(z) - m(a))$, so $f(z)$ can be written as $m(a) \log(\gamma, z)$ plus a function continuously extending to the analytic boundary for $z \sim \gamma(a)$. (i) follows by cutting γ into pieces. $\square$

A convenient lemma in the other direction is:

Lemma 2.3. *Suppose that $m \in \mathcal{L}^1([a, b])$, and $f(z)$ is a function on $\overline{\mathbb{C}} \setminus \Gamma$ such that*

- a.) *f is holomorphic on $\mathbb{C} \setminus \Gamma$, and f vanishes at ∞ ;*
- b.) *f extends to a distribution on $\mathbb{C}$;*
- c.) *if $s \in (a, b)$ then there are functions f^+ and f^- coming from continuous extensions of f to the analytic boundary on the left and right side of γ respectively, and*

$$m(t) = \frac{f^+(z) + f^-(z)}{2\pi i};$$

d.) for $z=\gamma(a)$ and $z=\gamma(b)$ there exists a direction η which is not a curve direction at z and

$$\lim_{\varepsilon \searrow 0} \varepsilon f(z + \varepsilon \eta) = 0.$$

Then, we claim,

$$(2) \quad f(z) = \int m(t) \frac{d\gamma(t)}{\gamma(t) - z}.$$

Proof. Let temporarily g denote the right side of (2). If g is such a function then due to 2.2.f the distribution $\bar{\partial}(g - f)$ is a distribution supported at the endpoints of γ . Then consider $r = (z \mapsto \frac{1}{\pi z}) * \bar{\partial}(g - f)$. We see that r is a rational function with singular support at the endpoints of γ at most, and one which vanishes at ∞ . Also, $\bar{\partial}$ vanishes on $(g - f) - r$. Then, from Liouville's theorem we see that $g - f = r$. Due to 2.2.e and condition (d) here we see that $g - f$ has actually no singular support, consequently $g - f = 0$. $\square$

2.4. As an immediate consequence of this lemma we may deduce:

a.) For $n \in \mathbb{N}$

$$\log(\gamma, z)^{n+1} = \int \frac{1}{2\pi i} ((\log(\gamma, \gamma(t)) + \pi i)^{n+1} - (\log(\gamma, \gamma(t)) - \pi i)^{n+1}) \frac{d\gamma(t)}{\gamma(t) - z}.$$

b.) For $|\alpha| < 1$

$$\exp \alpha \log(\gamma, z) = 1 + \int \frac{\sin \alpha \pi}{\pi} \exp \alpha \log(\gamma, \gamma(t)) \frac{d\gamma(t)}{\gamma(t) - z}.$$

2.5. In the special case when $[a, b] = [-1, 1]$ and $\gamma(t) = t$ it yields

a.) For $n \in \mathbb{N}$

$$\left(\log \left(\frac{z-1}{z+1} \right) \right)^{n+1} = \int \frac{1}{2\pi i} \left(\left(\log \left(\frac{1-t}{1+t} \right) + \pi i \right)^{n+1} - \left(\log \left(\frac{1-t}{1+t} \right) - \pi i \right)^{n+1} \right) \frac{dt}{t-z}.$$

b.) For $|\alpha| < 1$

$$\left(\frac{z-1}{z+1} \right)^\alpha = 1 + \int \frac{\sin \alpha \pi}{\pi} \left(\frac{1-t}{1+t} \right)^\alpha \frac{dt}{t-z}.$$

One can make similar statements concerning Lipschitz curve systems. There the role of endpoints will be played by the nodes of the curve system.

3. CERTAIN ESTIMATES AND APPLICATIONS TO CAUCHY INTEGRALS

A. First, we prove some estimates.

3.1. Suppose that $\mu \geq 0$ is a measure on the set S and $f : S \rightarrow [0, +\infty]$ is a measurable function. Then we define the straightening of f along μ as the function $\langle f | \mu \rangle : [0, \infty) \rightarrow [0, +\infty]$ given by

$$\langle f | \mu \rangle(x) = \inf \{ y : \mu(f > y) \leq x \}.$$

Simply speaking, the straightening operation rearranges the values of f into a monotone decreasing function meanwhile it takes into respect that “how much a given value occurs” according to μ . If $\mu(f > c) = +\infty$, then the value c is discarded. Using step-functions it is easy to prove

Lemma 3.2. *Let $\mu \geq 0$ be a measure, and $f_1, \dots, f_p$ be functions as above.*

a.) *Then*

$$\int_S f_1(t) \dots f_p(t) \mu(t) \leq \int_{[0, \infty)} \langle f_1 | \mu \rangle(x) \dots \langle f_p | \mu \rangle(x) dx.$$

b.) *If the pairing is monotone in the sense that for all $1 \leq i, j \leq p$ and $t_1, t_2 \in S$ $f_i(t_1) < f_i(t_2)$ implies $f_j(t_1) \leq f_j(t_2)$ then we have equality.* $\square$

3.3. We introduce some terminology, so we can formulate some technical statements about Lipschitz curves more easily. If $\gamma : [a, b] \rightarrow \mathbb{C}$ is a short Lipschitz curve then we define the sets

$$\mathbf{C}_{\gamma(a)\gamma} = \left\{ \frac{\gamma(t_2) - \gamma(a)}{|\gamma(t_2) - \gamma(a)|} : t_1 > a \right\} \subset \mathbf{A}\gamma = \left\{ \frac{\gamma(t_2) - \gamma(t_1)}{|\gamma(t_2) - \gamma(t_1)|} : t_2 > t_1 \right\}$$

as the set of global curve directions at $\gamma(a)$ and the set of forward directions of γ , respectively. We can consider the closed cone of curve directions $\mathbb{R}^+ \mathbf{C}_{\gamma(a)\gamma}$ and the cone of forward directions $\mathbb{R}^+ \mathbf{A}\gamma$. Let us denote the semi-angle of the cone $\mathbb{R}^+ \mathbf{A}\gamma$ by α . This is always smaller than $\frac{\pi}{2}$ for such a short Lipschitz curve. One can see that $\bigcap_{t>a} \mathbf{C}_{\gamma(a)\gamma}|_{[a,t]}$ is the set of curve directions at $\gamma(a)$. For $s > 0$ we use the notation $\log^+ s = \max(0, \log s)$.

Lemma 3.4. *Suppose that γ is a short Lipschitz curve as above, $\xi \in \mathbb{C} \setminus (\mathbb{R}^+ \mathbf{C}_{\gamma(a)\gamma})$. Suppose that the (minimal) angle between the direction of ξ and those of $\mathbf{C}_{\gamma(a)\gamma}$ is β .*

a.) *Let $f(t) = \frac{1}{|\gamma(t) - (\gamma(a) + \xi)|}$, $\mu(t) = |d\gamma(t)|$. Let ρ be the angle between the direction of ξ and the axis of $\mathbb{R}^+ \mathbf{A}\gamma$. Let $\alpha_1 = \rho - \beta$. Notice that $-\alpha \leq \alpha_1 \leq \alpha$. Let us set $\psi(t) = \min(t \tan \alpha_1, |\xi| \sin(\alpha_1 + \beta))$ for $t \in [0, \infty]$ and $F(t) = \frac{1}{|\xi|e^{i(\alpha_1 + \beta)} - (t + i\psi(t))|}$. Let $\tilde{\lambda}$ be $\frac{1}{\cos \alpha}$ times the Lebesgue measure restricted to $[0, \infty)$. Then $\langle f | \mu \rangle(x) \leq \langle F | \tilde{\lambda} \rangle(x)$.*

b.) *We claim that*


$$\int \frac{1}{|\gamma(t) - (\gamma(a) + \xi)|^2} |d\gamma(t)| \leq \frac{\pi}{|\xi| \beta \cos \alpha}.$$

c.) *If the arc-length of γ is ℓ then*

$$\int \frac{1}{|\gamma(t) - (\gamma(a) + \xi)|} |d\gamma(t)| \leq \frac{2}{\cos \alpha} \left(1 + \log^+ \frac{\ell \pi}{2|\xi| \beta} \right).$$

The latter two inequalities are monotone in α and β , so they apply if the semi-angle of $\mathbb{R}^+ \mathbf{A}\gamma$ is at most α , and the angle between the direction of ξ and those of $\mathbf{C}_{\gamma(a)\gamma}$ is at least β .

Proof. a.) We can make the assumptions that $\gamma(a) = 0$, the axis of $\mathbb{R}^+ \mathbf{A}\gamma$ is the positive x -axis, $\operatorname{Re} \xi \geq 0$, and γ is parametrized by its x coordinate.



b.) $\int_{[a,b]} |f(t)|^2 |d\gamma(t)| \leq \int_{[0,\infty)} |F(x)|^2 \frac{dx}{\cos \alpha}$, and the latter term can be computed exactly. If $\alpha_1 > 0$ then the latter integral is composed of two pieces and integration yields

$$= \frac{(\pi - \alpha_1 - \beta) \cos \alpha_1 + \sin \alpha_1}{|\xi| \sin \beta \cos \alpha} \leq \frac{(\pi - \alpha_1 - \beta) + \sin \alpha_1}{|\xi| \sin \beta \cos \alpha} \leq \frac{\pi - \beta}{|\xi| \sin \beta \cos \alpha} \leq \frac{\pi}{|\xi| \beta \cos \alpha}.$$

If $\alpha_1 < 0$ then there is one piece and it yields $\frac{(\pi-\beta) \cos \alpha_1}{|\ell| \sin \beta \cos \alpha}$, which can be estimated alike.

c.) Let $G_\omega(x) = \frac{1}{\cos \alpha_1} \frac{1}{|e^{i(\alpha_1 + \omega)} - x(1 + i \tan \alpha_1)|}$. Let $\hat{\beta} = \min(\beta, \frac{\pi}{2})$. Then one can see that $F(x) \leq G_\beta(x) \leq G_{\hat{\beta}}(x)$ if $x \geq 0$. Consequently,

$$\int \frac{1}{|\gamma(t) - (\gamma(a) + \xi)|} |d\gamma(t)| \leq \int F(t) |d\gamma(t)| \leq \int G_{\hat{\beta}}(t) |d\gamma(t)|.$$

We estimate the last term. Doing that we can forget about the condition $t \geq 0$. Then the integral is maximal when $|d\gamma(t)|$ is concentrated around $|\xi| \cos \beta \cos \alpha_1$, but the density relative to the Lebesgue measure is at most $\frac{1}{\cos \alpha}$. Hence we can estimate as

$$\leq \int_{|\xi| \cos \beta \cos \alpha_1 - \frac{1}{2} \ell \cos \alpha}^{|\xi| \cos \beta \cos \alpha_1 + \frac{1}{2} \ell \cos \alpha} G_{\hat{\beta}}(t) \frac{dt}{\cos \alpha} = \frac{2}{\cos \alpha} \operatorname{ar sh} \frac{\ell \cos \alpha}{2|\xi| \sin \hat{\beta} \cos \alpha_1}.$$

Using $|\alpha_1| \leq \alpha$ and $\frac{\beta}{\pi} \leq \sin \hat{\beta}$ we can continue by

$$\leq \frac{2}{\cos \alpha} \operatorname{ar sh} \frac{\pi \ell}{2|\xi|\beta} \leq \frac{2}{\cos \alpha} \left(1 + \log^+ \frac{\pi \ell}{2|\xi|\beta} \right).$$

Lemma 3.5. *Suppose that γ is a short Lipschitz curve and ξ is a transversal vector in angle $\geq \theta$. Let $\varepsilon, c > 0$, $t_0 \in [a, b]$. Set $s = \gamma(t_0) + \varepsilon\xi$, $r(s) = \gamma(t_0) - c\varepsilon\xi$.*

Then, we claim,

$$\int_{[a,b]} \frac{1}{|\gamma(t) - s|} \frac{1}{|\gamma(t) - r(s)|} |d\gamma(t)| \leq \frac{2\pi(c^{1/2} + c^{-1/2})}{|r(s) - s| \sin^2 \theta}.$$

Proof. We cut the interval $[a, b]$ into pieces $[a, t_0]$ and $[t_0, b]$. For each piece (with possible change of orientation) point (b) of the previous lemma applies with the choice $\alpha = \frac{\pi}{2} - \theta$, $\beta = \theta$. Adding up the estimates on the pieces and applying $\sin \theta \leq \theta$ we obtain

$$\int_{[a,b]} \frac{1}{|\gamma(t) - s|^2} |d\gamma(t)| \leq \frac{2\pi}{\varepsilon \sin^2 \theta}; \quad \int_{[a,b]} \frac{1}{|\gamma(t) - r(s)|^2} |d\gamma(t)| \leq \frac{2\pi}{c\varepsilon \sin^2 \theta}.$$

Applying the Cauchy inequality we obtain the estimate $\leq \frac{2\pi}{\varepsilon\sqrt{c}\sin^2\theta}$. Substituting $\varepsilon = \frac{1}{1+c} |r(s) - s|$ the statement follows. $\square$

3.6. Now we generalize the statements above to Lipschitz curve systems. If $\Gamma \subset \mathbb{C}$ is a Lipschitz curve system, $w \in \Gamma$ then we can imagine the constituent curves of Γ as follows: We have short Lipschitz curves $\gamma : [a_i, b_i] \rightarrow \mathbb{C}$ ($1 \leq i \leq p$) such that $\gamma_i(a_i) = w$, and the sets $\mathbf{A}\gamma_i$ are pairwise disjoint. We say that these curves form a star system around w . Apart from these there may be other constituents of Γ but these do not affect the local structure around w . We often consider star systems as above for the purposes of local analysis. We generally denote the maximal cone semi-angle of the $\mathbb{R}^+\mathbf{A}\gamma_i$ by α , and we set $\mathbf{C}_w = \bigcup_{i=1}^p \mathbf{C}_w\gamma_i$.

Lemma 3.7. *Suppose that we have a star system around w as above. Assume that C is an open cone from $\mathbb{C} \setminus \{0\}$. Also assume that the angle between any vector in C and any vector in $\mathbf{C}_w$ is $\geq \beta$. (See Figure A.) Then, we claim, for any $\xi_1, \xi_2 \in C$*

$$\sum_{i=1}^p \int \frac{1}{|\gamma_i(t) - (w + \xi_1)| |\gamma_i(t) - (w + \xi_2)|} |d\gamma_i(t)| \leq \frac{\pi p}{\sqrt{|\xi_1| |\xi_2|} \beta \cos \alpha}.$$

Proof. This is immediate from Lemma 3.4.b and the Cauchy inequality. $\square$




Figure A.




Figure B.

3.8. Consider a star system as earlier. Let $p \geq 2$, so that we can formulate the statement in simpler terms. We can assume that the γ_i are listed such that sets $\mathbf{A}\gamma_i$ come in counterclockwise direction. Suppose that u^+, u^- are unit vectors such that u^+ points between $\mathbf{C}_w\gamma_1$ and $\mathbf{C}_w\gamma_2$, and u^- points between $\mathbf{C}_w\gamma_1$ and $\mathbf{C}_w\gamma_p$. In other words $\mathbb{R}^+u^+$ and $\mathbb{R}^+u^-$ close γ_1 together from left and right. We can assume that the angle between $\mathbb{R}^+\mathbf{C}_w\gamma_2$ and $\mathbb{R}^+u^+$ is $\geq \beta$, the angle between $\mathbb{R}^+u^+$ and $\mathbb{R}^+\mathbf{C}_w\gamma_1$ is β^+ , the angle between $\mathbb{R}^+\mathbf{C}_w\gamma_1$ and $\mathbb{R}^+u^-$ is β^- , the angle between $\mathbb{R}^+u^-$ and $\mathbb{R}^+\mathbf{C}_w\gamma_p$ is $\geq \beta$. We also assume that u^+ and u^- are in angle $\leq \phi < \frac{\pi}{2}$ from the axis of $\mathbb{R}^+\mathbf{A}\gamma_i$. We define the “sector” U^+ as the open set between the curve γ_1 and $\mathbb{R}^+u^+$, and the “sector” U^- as the open set between the curve γ_1 and $\mathbb{R}^+u^-$. These sets are not completely well-defined, but they are quite well-defined in a sufficiently small neighborhood of w , ie. locally. So,

we just use U^+ and U^- without further ado, although what follows works only locally. (See Figure B.)

Now, we can define a sort of reflection map $r : U^+ \rightarrow U^-$ as follows: Take a point $s \in U^+$, then draw a line perpendicular to the axis of $\mathbf{A}\gamma_1$. Suppose that this line intersects γ_1 at s_0 . Then we set $r(s) = s_0 + c(s_0 - s)$, where c is a positive constant. One can show that if $c < \frac{2 \tan(\beta^-/2)}{\tan \phi - \tan(\beta^+ - \phi)}$ then the map indeed goes $U^+ \rightarrow U^-$. The very crude estimate $|s - r(s)| \leq 2 \tan \phi \sqrt{|s| |r(s)|}$ also holds.

Lemma 3.9.

$$\sum_{i=1}^p \int \frac{1}{|\gamma_i(t) - s| |\gamma_i(t) - r(s)|} |d\gamma_i(t)| \leq \frac{G}{|r(s) - s|},$$

where

$$G = \frac{2\pi(c^{1/2} + c^{-1/2})}{\cos^2 \alpha} + \frac{2\pi(p-1) \tan \phi}{\beta \cos \alpha}.$$

Proof. From Lemma 3.5 we can estimate the term $i = 1$, from Lemma 3.7 we can estimate the terms $i > 1$ directly. $\square$

Regarding to the situation above we say that U^+ is a sector siding γ_1 from the right, and U^- is a sector siding γ_1 from the left. One can notice that if U^+ is siding γ_1 then so do the sectors obtained by closing u^+ to $\mathbb{R}^+ \mathbf{C}_w \gamma_1$, and similarly for U^- .

Lemma 3.10. *Suppose that γ is a short Lipschitz curve. Let $w \in \mathbb{C}$ be arbitrary, $c \geq 0$. Let $f(t) = \log^+ \frac{1}{|\gamma(t) - w|}$, $\mu(t) = |d\gamma(t)|$, here $S = \gamma([a, b])$. Suppose that ℓ is the arc-length of the curve. Then, we claim,*

$$\langle f | \mu \rangle(x) \leq \chi_{[0, \ell)}(x) \log^+ \frac{2}{x \cos \alpha}.$$

Proof. We can assume that γ is in arc-length parametrization, $[a, b] = [-\ell/2, \ell/2]$, the axis of $\mathbb{R}^+ \mathbf{A}\gamma$ is the x axis, $\operatorname{Re} \gamma(t)$ is monotone increasing. Let $s = t_0$ if $\operatorname{Re} \gamma(t_0) = \operatorname{Re} w$, $s = \ell/2$ if $\operatorname{Re} \gamma(\ell/2) < \operatorname{Re} w$, $s = -\ell/2$ if $\operatorname{Re} \gamma(-\ell/2) > \operatorname{Re} w$. Then $|\gamma(t) - w| \geq |\operatorname{Re} \gamma(t) - \operatorname{Re} w| \geq \cos \alpha |t - s|$. In particular, it is enough to consider $F(t) = \log^+ \frac{1}{|t - s| \cos \alpha}$ instead of $f(t)$ on $[-\ell/2, \ell/2]$. The straightening of F is maximal when $s = 0$, and it yields the formula in the statement. $\square$

Lemma 3.11. *Suppose that γ is a short Lipschitz curve. Suppose that ℓ is the arc-length of γ . Let $w_1, \dots, w_n \in \mathbb{C}$, and $c > 0$ be arbitrary. Then*

$$\begin{aligned} \int \left(c + \left| \log \frac{1}{|\gamma(t) - w_1|} \right| \right) \dots \left(c + \left| \log \frac{1}{|\gamma(t) - w_n|} \right| \right) |d\gamma(t)| &\leq \\ &\leq n! \ell \left(1 + c + \log^+ \frac{2}{\ell \cos \alpha} + \log^+ H \right)^n \end{aligned}$$

holds, where $H = \max\{|\gamma(t) - w_i| : 1 \leq i \leq n, t \in [a, b]\}$.

Proof. If $\{i_1, \dots, i_p\}$ is a subset of $\{1, \dots, n\}$ then

$$\begin{aligned} \int \log^+ \frac{1}{|\gamma(t) - w_{i_1}|} \dots \log^+ \frac{1}{|\gamma(t) - w_{i_p}|} |d\gamma(t)| &\leq \int \left(\chi_{[0, \ell)}(x) \log^+ \frac{2}{x \cos \alpha} \right)^p dx \\ (3) \quad &\leq \left[p! u \sum_{j=0}^p \frac{1}{j!} \left(\log^+ \frac{2}{u \cos \alpha} \right)^j \right]_{u=0}^{\ell} \leq n! \ell \left(1 + \log^+ \frac{2}{\ell \cos \alpha} \right)^p. \end{aligned}$$

Using $c + \left| \log \frac{1}{|\gamma(t) - w_i|} \right| \leq c + \log^+ \frac{1}{|\gamma(t) - w_i|} + \log^+ H$ we can decompose our original integral, and from (3) we obtain the estimate in the statement. $\square$

Lemma 3.12. *Suppose that $\Gamma \subset \mathbb{C}$ is a piecewise Lipschitz curve system, $n \in \mathbb{N}$. Then, we claim, there exist a number $C_{\Gamma, n} > 0$ such that for any Lipschitz curve γ lying in Γ and $w_1, \dots, w_n \in \mathbb{C}$, $c \geq 0$ arbitrary the inequality*

$$\begin{aligned} \int \left(c + \left| \log \frac{1}{|\gamma(t) - w_1|} \right| \right) \dots \left(c + \left| \log \frac{1}{|\gamma(t) - w_n|} \right| \right) |d\gamma(t)| \leq \\ \leq C_{\Gamma, n} \ell \left(1 + c + \log^+ \frac{1}{\ell} + \log^+ H \right)^n \end{aligned}$$

holds, where $H = \max\{|\gamma(t) - w_i| : 1 \leq i \leq n, t \in [a, b]\}$, and ℓ is the arc-length of γ .

Proof. This follows from the previous lemma by cutting Γ to small pieces. $\square$

3.13. One can similarly estimate

$$\int (c + |\gamma(t) - w_1|^{-\alpha_1}) \dots (c + |\gamma(t) - w_n|^{-\alpha_n}) |d\gamma(t)|$$

where $0 < \alpha_i$, $\sum_i \alpha_i < 1$ and many other functions. The concrete examples above are also easy $\mathcal{L}^p$ -estimates.

B. Applications to integrals of Cauchy type.

Proposition 3.14. *Consider the situation of 2.2.*

j.) *Assume that $(a_1, b_1) \subset [a, b]$ and $\mu|_{(a_1, b_1)}(t) = m(t) d\gamma(t)$ where m is continuous on (a_1, b_1) . Then we claim: If f extends to $\gamma((a_1, b_1))^-$ continuously then it also extends to $\gamma((a_1, b_1))^+$ continuously and vice versa. In particular, $(\dagger)$ holds on (a_1, b_1) .*

k.) *Suppose that on $[a, b_1) \subset [a, b]$ and $\mu|_{[a, b_1)}(t) = m(t) \gamma(t)$, where $m(t)$ is continuous. If η stays in a closed sector not containing any curve direction of γ at $\gamma(a)$ then*

$$\lim_{\eta \rightarrow 0} \frac{f(\gamma(a) + \eta)}{\log |\eta|} = -m(a).$$

Proof. j.) Due to the local compactness of $\gamma([a, b])^+$ it is enough to show that the infinitesimal oscillation of f near $\gamma(s)^+$ ($s \in (a, b)$) is 0. Suppose that the direction ξ is transversal to γ in angle $\geq \theta$ in a neighborhood of $\gamma(s)$. Suppose that $[a_2, b_2]$ is an even smaller neighborhood of s such that $\max_{t \in [a_2, b_2]} |m(s) - m(t)| < \delta$. We see that

$$\begin{aligned} f(\gamma(s) + u\xi) &= f(\gamma(s) - u\xi) + m(s) \log(\gamma, \gamma(s) + u\xi) - m(s) \log(\gamma, \gamma(s) - u\xi) \\ &+ \int_{[a, b] \setminus [a_2, b_2]} \left(\frac{1}{\gamma(t) - \gamma(s) - u\xi} - \frac{1}{\gamma(t) - \gamma(s) + u\xi} \right) (m(t) - m(s)) d\gamma(t) \\ &+ \int_{[a_2, b_2]} \left(\frac{1}{\gamma(t) - \gamma(s) - u\xi} - \frac{1}{\gamma(t) - \gamma(s) + u\xi} \right) (m(t) - m(s)) d\gamma(t). \end{aligned}$$

After the equality, as a function of $\gamma(s) + u\xi$, every term extends to $\gamma([a_2, b_2])^+$ continuously, except the last one, but which, according to Lemma 3.5, we can estimate by $\frac{2\pi}{\sin^2 \theta} \delta$. This shows that the infinitesimal oscillation of f near $\gamma(s)^+$ is at most $\frac{4\pi}{\sin^2 \theta} \delta$. The δ can be arbitrarily small, so we see that the infinitesimal oscillation of f near $\gamma(s)^+$ is actually 0.

k.) Suppose that $[a, b_1] \subset [a, b]$ is so small that $\gamma|_{[a, b_1]}$ is short and the semi-angle of $\mathbf{A}\gamma|_{[a, b_1]}$ is α . Let $[a, b_2] \subset [a, b]$ be even smaller so that $\max_{t \in [a, b_2]} |\gamma(t) - \gamma(a)| < \delta$.

$$\begin{aligned} f(\gamma(s) + \eta) &= m(a) \log(\gamma, \gamma(a) + \eta) + \int_{[a, b] \setminus [a, b_2]} (m(t) - m(a)) \frac{d\gamma(t)}{\gamma(t) - (\gamma(a) + \theta)} \\ &\quad + \int_{[a, b_2]} (m(t) - m(a)) \frac{d\gamma(t)}{\gamma(t) - (\gamma(a) + \eta)}. \end{aligned}$$

Dividing by $\log |\eta|$ we see that in the right side of the equation the first term has the required asymptotic behaviour, the second term goes to zero as we divide a bounded quantity by a logarithmic one. According to Lemma 3.4.c, the limsup of the absolute value of the first term divided by $\log |\eta|$ is less or equal to $\frac{2}{\cos \alpha} \delta$. Choosing δ well, this latter term can be arbitrarily small, so the statement follows. $\square$

3.15. Let us consider a star system Γ as in point 3.6. Assume that $m_j : [a_j, b_j] \rightarrow \mathbb{C}$ are continuous functions. For $z \in \mathbb{C} \setminus \Gamma$ let us define

$$f(z) = \sum_{j=1}^p \int m_j(t) \frac{d\gamma_j(t)}{\gamma_j(t) - z}.$$

Proposition 3.16. *Suppose that C is an open cone from $\mathbb{C} \setminus \{0\}$. Assume the angle between any two directions from C and $\mathbf{A}_w$ is $\geq \beta$. Suppose that $w_1 \notin w + \mathbb{R}^+ \mathbf{A}$, and $f(z)$ is bounded on the open segment T connecting w and w_1 . Then, we claim,*

a.) $\sum_{j=1}^p m(a_i) = 0$.

b.) $f(z)$ extends continuously to the analytic boundary of $w + C$.

Proof. a.) Let $\eta = w_1 - w$. According to Proposition 3.14.k $\lim_{\varepsilon \searrow 0} \frac{f(w + \varepsilon \eta)}{\log \varepsilon |\eta|} = - \sum_{j=0}^p m(a_i)$,

but which has to vanish according to the boundedness of f in the expression.

b.) The statement is already known except for around w . We can assume that $w = 0$. We can assume that C is so small that the angle between any two vectors in C is $\leq \frac{\pi}{2}$. Let L be a linear map sending C into $\mathbb{R}^+ w_1 \setminus \{0\}$. Choosing it well, we can assume that there is a positive number c such that for $z \in C$ we have $|z| \leq c|L(z)|$, $|L(z)| \leq c|z|$. Let us chose $\tilde{b}_i$ such that $a_i < \tilde{b}_i < b_i$ and

$$\max_{1 \leq i \leq p, t \in [a_i, \tilde{b}_i]} |m_i(t) - m_i(a_i)| < \delta.$$

Now, for $z \in C$

$$\begin{aligned} f(z) &= f(L(z)) + \sum_{i=1}^p m_i(a_i) \log(\gamma_i, z) - \sum_{i=1}^p m_i(a_i) \log(\gamma_i, L(z)) + \\ &\quad + \sum_{i=1}^p \int_{[\tilde{b}_i, b_i]} (m_i(t) - m_i(a_i)) \left(\frac{1}{\gamma_i(t) - z} - \frac{1}{\gamma_i(t) - L(z)} \right) d\gamma_i(t) + \\ &\quad + \int_{[a_i, \tilde{b}_i]} (m_i(t) - m_i(a_i)) \left(\frac{1}{\gamma_i(t) - z} - \frac{1}{\gamma_i(t) - L(z)} \right) d\gamma_i(t). \end{aligned}$$

Then, on the right side of the equation every term is continuous on C except possibly the last one. According to Lemma 3.7, the absolute value of the last term

$$= \left| \int_{[a_i, \tilde{b}_i]} (m_i(t) - m_i(a_i)) \frac{L(z) - z}{(\gamma_i(t) - z)(\gamma_i(t) - L(z))} d\gamma_i(t) \right| \leq$$

$$\leq \delta |L(z) - z| \frac{\pi p}{\sqrt{|z|} |L(z)| \beta \cos \alpha} \leq \frac{\pi p (c^{1/2} + c^{-1/2})}{\beta \cos \alpha} \delta.$$

In particular, the infinitesimal oscillation of f on C around w is less or equal than twice the value above. Now, δ can be arbitrarily small, so we know that f extends continuously. $\square$

Proposition 3.17. *Suppose that the sectors U^+ and U^- are siding γ_1 from left and right respectively. If f extends to the analytic boundary of U^- continuously then it also extends to the analytic boundary of U^+ continuously.*

Proof. It is similar to the previous ones, but we have to use Lemma 3.9. $\square$

4. RATIONAL FUNCTIONS AND RELATED IDENTITIES

Here we collect some identities of rational functions on $\mathbb{C}$.

Lemma 4.1. *Suppose that w_1, w_2, z are three different points in $\mathbb{C}$. Then*

a.)

$$\frac{1}{w_1 - z} \frac{1}{w_2 - z} = \frac{1}{w_2 - w_1} \frac{1}{w_1 - z} + \frac{1}{w_1 - w_2} \frac{1}{w_2 - z}.$$

b.) For $n \in \mathbb{N}$,

$$\frac{1}{w_1 - w_2} \left(\frac{1}{(w_1 - z)^n} - \frac{1}{(w_2 - z)^n} \right) = - \sum_{j=1}^n \frac{1}{(w_1 - z)^j} \frac{1}{(w_2 - z)^{n-j+1}}.$$

c.) For $n_1, n_2 \in \mathbb{N}$

$$\begin{aligned} \frac{1}{(w_1 - z)^{n_1+1} (w_2 - z)^{n_2+1}} &= \sum_{j=0}^{n_1} \frac{(j + n_2)!}{j! n_2!} \frac{1}{(w_2 - w_1)^{n_2+j+1} (w_1 - z)^{n_1-j+1}} + \\ &+ \sum_{j=0}^{n_2} \frac{(j + n_1)!}{j! n_1!} \frac{1}{(w_1 - w_2)^{n_1+j+1} (w_2 - z)^{n_2-j+1}}. \end{aligned}$$

Now assume that $\gamma : [a, b] \rightarrow \mathbb{C}$ is a Lipschitz curve, $\Gamma = \gamma([a, b])$. Drawing on the basic integral formula we can get finer statements. In what follows $z \in \mathbb{C} \setminus \Gamma$.

Lemma 4.2. a.) *Suppose that $a_1, b_1 \in [a, b]$ and $n \geq 1$. Then*

$$\frac{1}{(\gamma(b_1) - z)^n} - \frac{1}{(\gamma(a_1) - z)^n} = \int_{t=a_1}^{b_1} \frac{-n d\gamma(t)}{(\gamma(t) - z)^{n+1}}.$$

b.) *Suppose that $a_1, b_1 \in [a, b]$ and*

$$\sum_{j=1}^n \frac{c_j}{(-w)^j} \sim \sum_{j=1}^n \frac{c_j^{(\gamma(a_1))}}{(\gamma(a_1) - w)^j} \sim \sum_{j=1}^n \frac{c_j^{(\gamma(b_1))}}{(\gamma(b_1) - w)^j}$$

modulo $O(w^{-n-1})$ as $w \rightarrow \infty$. Then

$$\sum_{j=1}^n \frac{c_j^{(\gamma(b_1))}}{(\gamma(b_1) - z)^j} - \sum_{j=1}^n \frac{c_j^{(\gamma(a_1))}}{(\gamma(a_1) - z)^j} = \int_{t=a_1}^{b_1} \left(\sum_{j=1}^n \binom{n-1}{j-1} c_j \gamma(t)^{n-j} \right) \frac{-n d\gamma(t)}{(\gamma(t) - z)^{n+1}}.$$

c.) Suppose that $a \leq a_1 < b_1 \leq b$ and $n_1, n_2 \in \mathbb{N}$. Then

$$\begin{aligned} & \frac{1}{(\gamma(a_1) - z)^{n_1+1}(\gamma(b_1) - z)^{n_2+1}} = \\ &= \int_{t=a_1}^{b_1} \frac{(n_1 + n_2 + 1)!}{n_1!n_2!} \frac{(\gamma(t) - \gamma(a_1))^{n_2}(\gamma(b_1) - \gamma(t))^{n_1}}{(\gamma(b_1) - \gamma(a_1))^{n_1+n_2+1}} \frac{d\gamma(t)}{(\gamma(t) - z)^{n_1+n_2+2}}. \end{aligned}$$

Next, we are going to examine superpositions of the identities before.

4.3. Suppose that K and S are $\mathbf{T}_2, \mathbf{M}_2$, locally compact topological spaces, and we assign to each point $s \in S$ a measure ν_s on K . We say that this assignment is continuous if

- a.) the variation map $s \mapsto \|\nu_s\|$ is bounded and continuous;
- b.) the map $s \mapsto \nu_s$ is weakly continuous on S .

In this case, if μ is a measure on S then we may form the parametrized measure

$$\tilde{\theta}(t, s) = \nu_s(t)\mu(s),$$

on $S \times T$, most readily to be defined as a Radon measure. Integrating in the variable s we obtain the superposition measure

$$\theta(t) = \int_s \nu_s(t)\mu(s).$$

In fact, both the parametrized and the superposition measures can also be taken if $s \mapsto \nu_s$ is not necessarily continuous but S is a disjoint union of the Borel sets $S_1, \dots, S_p$ such that on each of them $s \mapsto \nu_s$ is continuous, and the S_j are $\mathbf{T}_2, \mathbf{M}_2$, locally compact spaces themselves.

Now, if $v(t)$ is a bounded continuous, locally convex vector space valued function then

$$(4) \quad \int \left(\int v(t) \nu_s(t) \right) \mu(s) = \int v(t) \tilde{\theta}(t, s) = \int v(t) \theta(t).$$

We will consider superpositions only in very special cases, where the general arguments above can also be carried out by more direct means.

4.4. We introduce the measure $\Omega_{a_1, b_1}^{(\gamma)}(t)$ on $[a, b]$. We define the function $\chi_{a, b}(x)$ such that $\chi_{a, b}(x) = \chi_{[a, b]}(x)$ if $a < b$, $\chi_{a, b}(x) = 0$ if $a = b$, and $\chi_{a, b}(x) = -\chi_{[b, a]}(x)$ if $b < a$. Then we set

$$\Omega_{a_1, b_1}^{(\gamma)}(t) = -\chi_{a_1, b_1}(t) d\gamma(t).$$

With this notation

$$\frac{1}{(\gamma(b_1) - z)^n} - \frac{1}{(\gamma(a_1) - z)^n} = \int \frac{n \Omega_{a_1, b_1}^{(\gamma)}(t)}{(\gamma(t) - z)^{n+1}}.$$

The map $(a_1, b_1) \mapsto \Omega_{a_1, b_1}^{(\gamma)}(t)$ is continuous in the sense of 4.3.

Lemma 4.5. Suppose that $\mu(t)$ is finite measure on $[a, b]$. Suppose that $t_0 \in [a, b]$, $n \in \mathbb{N}$. Then

$$\int \frac{\mu(t)}{(\gamma(t) - z)^n} = \frac{c_n}{(\gamma(t_0) - z)^n} + \int \frac{n \theta(t)}{(\gamma(t) - z)^{n+1}},$$

where

$$c_n = \int \mu(s) \quad \text{and} \quad \theta(t) = \int_s \Omega_{t_0, s}^{(\gamma)}(t) \mu(s).$$

$$\text{More explicitly, } \theta(t) = p(t) d\gamma(t) \text{ where } p(t) = \begin{cases} \int_{s=a}^t \mu(s) & \text{if } t < t_0 \\ -\int_{s=t}^b \mu(s) & \text{if } t > t_0. \end{cases}$$

4.6. Similarly to the additive case, let us define the measure $\Xi_{a_1, b_1}^{(\gamma, n_1, n_2)}(t)$ on $[a, b]$ as follows: If $a_1 < b_1$ then we set

$$\Xi_{a_1, b_1}^{(\gamma, n_1, n_2)}(t) = \chi_{[a_1, b_1]}(t) \frac{(n_1 + n_2 + 1)!}{n_1! n_2!} \frac{(\gamma(t) - \gamma(a_1))^{n_2} (\gamma(b_1) - \gamma(t))^{n_1}}{(\gamma(b_1) - \gamma(a_1))^{n_1 + n_2 + 1}} d\gamma(t);$$

if $a_1 = b_1$ then we set $\Xi_{a_1, b_1}^{(\gamma, n_1, n_2)}(t) = \delta_{a_1}(t)$, the one-point measure at $t = a_1$; and if $a_1 > b_1$ then we define $\Xi_{a_1, b_1}^{(\gamma, n_1, n_2)}(t) = \Xi_{b_1, a_1}^{(\gamma, n_1, n_2)}(t)$. Then

$$\frac{1}{(\gamma(a_1) - z)^{n_1 + 1} (\gamma(b_1) - z)^{n_2 + 1}} = \int \frac{\Xi_{a_1, b_1}^{(\gamma, n_1, n_2)}(t)}{(\gamma(t) - z)^{n_1 + n_2 + 2}}.$$

The map $(a_1, b_1) \mapsto \Xi_{a_1, b_1}^{(\gamma, n_1, n_2)}$ is continuous in the sense of 4.3 on the sets $\Delta^+ = \{(x, y) \in [a, b] \times [a, b] : x < y\}$, $\Delta = \{(x, y) \in [a, b] \times [a, b] : x = y\}$, and $\Delta^- = \{(x, y) \in [a, b] \times [a, b] : x > y\}$, separately. Indeed, only the boundedness property is nontrivial, but from the bi-Lipschitz property we obtain $\|\Xi_{a_1, b_1}^{(\gamma, n_1, n_2)}\| \leq (C_2/C_1)^{n_1 + n_2 + 1}$.

Lemma 4.7. Suppose that μ_1, μ_2 are finite measures on $[a, b]$, $n_1, n_2 \in \mathbb{N}$. Then

$$\int \frac{\mu(t)}{(\gamma(t) - z)^{n_1 + n_2 + 2}} = \int \frac{\mu_1(t_1)}{(\gamma(t_1) - z)^{n_1 + 1}} \int \frac{\mu_2(t_2)}{(\gamma(t_2) - z)^{n_2 + 1}}$$

holds with

$$\mu(t) = \int_{(s_1, s_2) \in [a, b] \times [a, b]} \Xi_{s_1, s_2}^{(\gamma, n_1, n_2)}(t) \mu_1(s_1) \mu_2(s_2),$$

ie. a superposition with respect to the product measure $\mu_1 \times \mu_2$.

4.8. The next step is to pass from Lipschitz curves to Lipschitz curve systems. In these cases, however, the normalization processes are not entirely canonical.

We start with the additive case. Suppose that $\Gamma \subset \mathbb{C}$ is a connected Lipschitz curve system, $s_0 \in \Gamma$. Then we can construct a choice function $\phi : \Gamma \rightarrow \Phi$, such that

- a.) Φ is a finite set of possibly degenerate Lipschitz curves $\gamma_i : [a_i, b_i] \rightarrow \mathbb{C}$ ($1 \leq i \leq p$) lying in Γ ;
- b.) if $\phi(s) = \gamma_i$ then $\{s_0, s\} \subset \gamma_i([a_i, b_i])$;
- c.) ϕ is a Borel function.

We define the measure $\Omega_{s_0, s}^{(\phi)}$ as $(\gamma_i)_* \Omega_{\gamma_i^{-1}(s_0), \gamma_i^{-1}(s)}^{(\gamma_i)}$ if $\phi(s) = \gamma_i$. The map $s \mapsto \Omega_{s_0, s}^{(\phi)}$ is such that we can consider superposition in the sense of 4.3:

Lemma 4.9 (Additive reduction). Suppose that $\Gamma \subset \mathbb{C}$ is a connected Lipschitz curve system, $w_0 \in \Gamma$, $n \in \mathbb{N}$, and the map $s \mapsto \Omega_{w_0, s}^{(\phi)}$ is constructed according to a suitable choice function ϕ . Suppose that μ is a measure on Γ . Then

$$\int \frac{\mu(w)}{(w - z)^n} = \frac{c_n}{(w_0 - z)^n} + \int \frac{n \theta(w)}{(w - z)^{n+1}}$$

where

$$c_n = \int \mu(s) \quad \text{and} \quad \theta(w) = \int_s \Omega_{w_0, s}^{(\phi)}(w) \mu(s).$$

If ϕ is piecewise constant on Γ then $\theta(w) = p(w)d\Gamma(w)$ piecewise, where $d\Gamma$ is the natural variation measure oriented in some direction and $w(s)$ is an integral function of $\mu(s)$ in the same direction.

4.10. We can deal with the multiplicative case as follows: Suppose that $\Gamma \subset \mathbb{C}$ is a Lipschitz curve system. Then we can construct a choice function $\phi : \Gamma \times \Gamma \rightarrow \Phi$, such that

a.) Φ is a finite set containing the empty set $\emptyset$ and possibly degenerate Lipschitz curves $\gamma_i : [a_i, b_i] \rightarrow \mathbb{C}$ ($1 \leq i \leq p$) lying in Γ ;

b₁.) if $\phi(s_1, s_2) = \gamma_i$ then $\{s_1, s_2\} \subset \gamma_i([a_i, b_i])$;

b₂.) there is constant $c > 0$ such that $\phi(s_1, s_2) = \emptyset$ implies $|s_1 - s_2| \geq c$;

c.) ϕ is a Borel function.

(The most natural case is when $\phi(s_1, s_2) = \emptyset$ if and only if s_1 and s_2 are in different connected components.)

We define the measures $\Xi_{s_1, s_2; k}^{(\phi, n_1, n_2)}$ ($1 \leq k \leq n_1 + n_2 + 2$) such that

$$\Xi_{s_1, s_2; k}^{(\phi, n_1, n_2)} = \begin{cases} \delta_{n_1+n_2+2, k}(\gamma_i) * \Xi_{\gamma_i^{-1}(s_1), \gamma_i^{-1}(s_2)}^{(\gamma_i, n_1, n_2)} & \text{if } \phi(s_1, s_2) = \gamma_i \\ \sum_{j=0}^{n_1} \frac{(j+n_2)!}{j!n_2!} \frac{\delta_{n_1-j+1, k} \delta_{s_1}}{(s_2 - s_1)^{n_2+j+1}} + \sum_{j=0}^{n_2} \frac{(j+n_1)!}{j!n_1!} \frac{\delta_{n_2-j+1, k} \delta_{s_2}}{(s_1 - s_2)^{n_1+j+1}} & \text{if } \phi(s_1, s_2) = \emptyset. \end{cases}$$

Then the maps $(s_1, s_2) \mapsto \Xi_{s_1, s_2; k}^{(\phi, n_1, n_2)}(s)$ are such that we can consider superposition in the sense of 4.3:

Lemma 4.11 (Multiplicative reduction). *Suppose that $\Gamma \subset \mathbb{C}$ is a Lipschitz curve system, μ_1 and μ_2 are measures on it. Suppose that the map $(s_1, s_2) \mapsto \Xi_{s_1, s_2; k}^{(\phi, n_1, n_2)}(s)$ is obtained according to a choice function ϕ as above. Then*

$$\int \frac{\mu_1(s_1)}{(s_1 - z)^{n_1+1}} \int \frac{\mu_2(s_2)}{(s_2 - z)^{n_2+1}} = \sum_{k=1}^{n_1+n_2+2} \int \frac{\theta_k(s)}{(s - z)^k},$$

where

$$\theta_k(s) = \int_{(s_1, s_2) \in \Gamma \times \Gamma} \Xi_{s_1, s_2; k}^{(\phi, n_1, n_2)}(s) \mu_1(s_1) \mu_2(s_2).$$

The following statements deal with global conformal transformations.

Lemma 4.12. *Let $h : \overline{\mathbb{C}} \rightarrow \overline{\mathbb{C}}$ be a conformal map, $h(z) = \frac{az+b}{cz+d}$, $ad - bc \neq 0$.*

a.) *If $w, h(w) \in \mathbb{C}$, $z \neq w$, $h(z) \in \mathbb{C}$ then*

$$\frac{1}{w - z} = \frac{c}{cw + d} + \frac{ad - bc}{(cw + d)^2} \frac{1}{h(w) - h(z)}.$$

b.) *Suppose that $\Gamma \subset \mathbb{C}$ is a Lipschitz curve system, μ is a measure on Γ . Assume that $h(\Gamma) \subset \mathbb{C}$. Suppose that $z \in \mathbb{C} \setminus \Gamma$ and $h(z) \in \mathbb{C}$. Then*

$$\int \frac{\mu(w)}{(w - z)^n} = \sum_{j=0}^n \int \binom{n}{j} \left(\frac{c}{cw + d} \right)^{n-j} \left(\frac{ad - cd}{(cw + d)^2} \right)^j \frac{\mu(w)}{(h(w) - h(z))^j}.$$

For the purpose of the following statements let $\mathcal{U} \subset \mathbb{C}$ be a simply connected Lipschitz domain with oriented boundary curve $\hat{\gamma}$ with underlying curve $\gamma : [a, b]_{a \sim b} \rightarrow \mathbb{C}$.

Lemma 4.13. *For any polynomial P with degree $\leq n - 1$*

$$\int P(\gamma(t)) \frac{d\gamma(t)}{(\gamma(t) - z)^{n+1}} = 0.$$

Lemma 4.14. *Assume that $f : \mathcal{U} \rightarrow \mathbb{C}$ is a holomorphic function which extends to the analytic boundary of $\mathcal{U}$ continuously. Suppose that $\hat{f} : [a, b]_{a \sim b} \rightarrow \mathbb{C}$ gives the boundary values in the sense that at $\hat{\gamma}(t)$ the extended value is $\hat{f}(t)$. Suppose that $n \in \mathbb{N}$, $z \in \mathbb{C} \setminus \overline{\mathcal{U}}$. Then*

$$\int \hat{f}(t) \frac{d\gamma(t)}{(\gamma(t) - z)^{n+1}} = 0.$$

Lemma 4.15. *Suppose that $n \geq 1$, $z \in \mathbb{C} \setminus \overline{\mathcal{U}}$, $t_0 \in [a, b] / \{a \sim b\}$.*

a.) If $w \in \mathcal{U}$ then

$$\frac{1}{(w - z)^n} = \frac{1}{(\gamma(t_0) - z)^n} + n \frac{1}{2\pi i} \int \log(\gamma(t) - w) \frac{d\gamma(t)}{(\gamma(t) - z)^{n+1}},$$

where $\log(\gamma(t) - w)$ is a branch such that it is continuous on $([a, b] / \{a \sim b\}) \setminus \{t_0\}$.

b.) If μ is a measure $\mathcal{U}$ and $z \in \mathbb{C} \setminus \overline{\mathcal{U}}$ then

$$\int \frac{\mu(w)}{(w - z)^n} = \frac{1}{(\gamma(t_0) - z)^n} \int \mu(w) + n \frac{1}{2\pi i} \int r(t, w) \frac{\mu(w) d\gamma(t)}{(\gamma(t) - z)^{n+1}}$$

assumed that i.) for a fixed w the function $r(t, w) = \log(\gamma(t) - w)$ is a branch such that it is continuous except at t_0 , ii.) $r(t, w) \mu(w) d\gamma_i(t)$ is Lebesgue integrable.

Such choice functions $r(t, w)$ can always be found.

5. NORMAL FORMS OF SLOWLY GROWING HOLOMORPHIC FUNCTIONS

5.1. Let $\overline{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ be the Riemann sphere. Suppose that $\Gamma \subset \overline{\mathbb{C}}$ is a closed set. We define $\mathcal{H}_\Gamma$ as the set of functions f such that a.) f is holomorphic on $\overline{\mathbb{C}} \setminus \Gamma$; b.) f is slowly growing around Γ , ie. there exist $n \in \mathbb{N}$, and $K > 0$ such that $|f(z)| \leq K \text{dist}_{\overline{\mathbb{C}}}(z, \Gamma)^{-n}$, where $\text{dist}_{\overline{\mathbb{C}}}$ is the distance function according to a Riemannian metric on $\overline{\mathbb{C}}$. The growth condition says that f grows around Γ polynomially with exponent at most n . (If $\Gamma \subset \mathbb{C}$ then this local condition can be checked using the ordinary Euclidean distance function dist .) It is easy to see that $\mathcal{H}_\Gamma$ is an algebra, the algebra of slowly growing holomorphic functions on $\overline{\mathbb{C}} \setminus \Gamma$. We will be interested primarily in the very special case when Γ is a Lipschitz curve system.

A. First we examine the case of Lipschitz curves on $\mathbb{C}$:

5.2. Let $\gamma : [a, b] \rightarrow \mathbb{C}$ be a Lipschitz curve, $\Gamma = \gamma([a, b])$. A function $f \in \mathcal{H}_\Gamma$ has a power series expansion around ∞ :

$$f(z) \sim c_0 + \frac{c_1}{-z} + \frac{c_2}{(-z)^2} + \frac{c_3}{(-z)^3} + \dots + \frac{c_n}{(-z)^n} + \dots$$

We define the operation of global primitive function M if $c_0 = c_1 = 0$. Then $Mf(z)$ is the primitive function which vanishes at ∞ :

$$Mf(z) \sim \frac{c_2}{-z} + \frac{c_3}{2(-z)^2} + \frac{c_4}{3(-z)^3} + \dots + \frac{c_{n+1}}{n(-z)^n} + \dots$$

It is easy to see that Mf is still in $\mathcal{H}_\Gamma$. It is often useful to write the expansion around infinity in terms of $z - p$, so

$$f(z) \sim c_0^{(p)} + \frac{c_1^{(p)}}{p - z} + \frac{c_2^{(p)}}{(p - z)^2} + \frac{c_3^{(p)}}{(p - z)^3} + \dots + \frac{c_n^{(p)}}{(p - z)^n} + \dots$$

We can notice that $c_0^{(p)} = c_0$ and $c_1^{(p)} = c_1$ but the other terms do vary. Nevertheless, $c_0^{(p_1)}, \dots, c_n^{(p_1)}$ can always be expressed from $c_0^{(p_2)}, \dots, c_n^{(p_2)}$. They encode the same information, the asymptotic expansion modulo $O(z^{-n-1})$.

5.3. Suppose that $n \in \mathbb{N}$, μ is a measure on $[a, b]$, such that

$$f(z) = \sum_{j=0}^n \frac{c_j^{(\gamma(t_0))}}{(\gamma(t) - z)^j} + \int \frac{\mu(t)}{(\gamma(t) - z)^{n+1}}.$$

In this case we say that $f(z)$ is written in a simple $(n+1)$ -normal form with base point $\gamma(t_0)$. A priori we do not ask that $c_j^{(\gamma(t_0))}$ should be understood in the sense of the previous point, but it is easy to see that they must be the same. It is reasonable to say that $f(z)$ allows a simple (0) -normal form if $f(z) = c_0$, independent of base points.

More generally, if $\mu_0, \dots, \mu_{n+1}$ are measures on $[a, b]$ and

$$f(z) = \sum_{j=0}^{n+1} \int \frac{\mu_k(t)}{(\gamma(t) - z)^k}$$

then we say that $f(z)$ is written in an $(n+1)$ -normal form.

On the other hand, we may also notice that if a function is of shape like above then it is a slowly growing holomorphic function, in fact, of exponent $n+1$.

5.4. A simple (n) -normal form is certainly an (n) -normal form. On the other hand, if $f(z)$ allows an (n) -normal form then it allows a simple (n) -normal form with any base point in $\gamma([a, b])$. Indeed, the measure μ_0 can be gathered into a constant c_0 , then by induction, using Lemma 4.5, we can push the measures up to the $(n+1)$ -th level, except the singular term in the base point. In fact, this gives a well-defined reduction process, whose validity is based solely on Lemma 4.5.

Proposition 5.5. *Every $f \in \mathcal{H}_{\gamma([a, b])}$ allows a simple (n) -normal form for certain n . If n and the base point are fixed then the simple normal form is unique.*

Proof. First we prove unicity. The $c_j^{\gamma(t_0)}$ are determined by the power series expansion of f around infinity. Then it is enough to show that $\int \frac{\mu(t)}{(\gamma(t) - z)^{n+1}}$ determines μ . Indeed, its n -th primitive function is $\frac{1}{n!} \int \frac{\mu(t)}{\gamma(t) - z}$, from which μ can be recovered according to the $\bar{\partial}$ -formula, 2.2.d.

Existence is as follows: If $f(z)$ is of exponent $n-2$ then locally its n -th primitive function extends Lipschitz continuously to the analytic boundary. However, there are obstructions to have a global primitive function. For that reason we take the global n -th primitive function $g(t)$ of $f(z) - \sum_{j=0}^n \frac{c_j^{\gamma(t_0)}}{(\gamma(t) - z)^j}$. We see this primitive function locally Lipschitz continuously extends to the analytic boundary except at $\gamma(t_0)^\pm$. Let $m(t) = \frac{1}{2\pi i} (g^+(z) - g^-(z))$. This is a locally Lipschitz function except at $\gamma(t_0)$. If we choose another $\tilde{t}_0$ and proceed with that then we see that the difference $\tilde{g}(z) - g(z)$ in the n -th primitive functions is $\frac{1}{n!} \int n \chi_{\gamma(\tilde{t}_0), \gamma(t_0)}(t) \left(\sum_{j=1}^n \binom{n-1}{j-1} c_j \gamma(t)^{n-j} \right) \frac{d\gamma(t)}{\gamma(t) - z}$. Now, m ($\tilde{m}$) can be recovered from g ($\tilde{g}$) using the $\bar{\partial}$ operator, and which yields $\tilde{m} - m = \frac{1}{n!} n \chi_{\gamma(\tilde{t}_0), \gamma(t_0)}(t) \left(\sum_{j=1}^n \binom{n-1}{j-1} c_j \gamma(t)^{n-j} \right)$. From that we can see that the discontinuity of $m(t)$ is not at all too bad, $m(t)$ is a piecewise Lipschitz function. Let $h = \sum_{j=0}^n \frac{c_j^{\gamma(t_0)}}{(\gamma(t) - z)^j} +$

$n! \int m(t) \frac{d\gamma(t)}{(\gamma(t)-z)^{n+1}}$. We see that $f - h$ extends holomorphically to $\overline{\mathbb{C}}$ except at $\gamma(t_0)$, and it is vanishing at infinity in all orders. From this we deduce that $f = h$. $\square$

B. Next, we assume that $\Gamma \subset \mathbb{C}$ is a Lipschitz curve system.

5.6. Suppose that Γ_j ($1 \leq j \leq b_0$) are the connected components of Γ . Let $w_h \in \Gamma_h$ be chosen. If with appropriate n , constants $c_j^{(w_h)}$, and a measure μ on Γ

$$f(z) = \sum_{h=1}^{b_0} \sum_{j=0}^n \frac{c_j^{(w_h)}}{(w_h - z)^j} + \int_{\Gamma} \frac{\mu(w)}{(w - z)^{n+1}}$$

then we say that $f(z)$ is written in a simple $(n+1)$ -normal form. Similarly, if $\mu_0, \dots, \mu_{n+1}$ are measures on Γ and

$$f(z) = \sum_{j=0}^{n+1} \int \frac{\mu_k(w)}{(w - z)^k}$$

then we say that $f(z)$ is written in an $(n+1)$ -normal form.

5.7. For Lipschitz curve systems things go in similar manner: (n) -normal forms can be reduced to simple (n) -normal forms by Lemma 4.9, every element f of $\mathcal{H}_{\Gamma}$ allows a normal form, etc. There is, however, one important difference: it is not true that (the w_h chosen) the simple normal form is unique. First, the constant terms $c_0^{(w_h)}$ can be accumulated into a single constant term. Apart from that, the higher $c_j^{(w_h)}$ are fixed, because they are obstructions to taking the j -th primitive functions around Γ_h . (In fact the Γ_h can be nested so that one encloses the other one but we can go in the inside-out direction.) But the μ can also be a source of ambiguity. Assume that $n > 1$ and Γ contains a cycle ζ . Assume that P is a polynomial of order $\leq n-1$. Then

$$\int \frac{P(\zeta) d\zeta}{(\zeta - z)^{n+1}} = 0.$$

Clearly, a term $P(\zeta) d\zeta$ can be added the measure-term μ without consequences. Let b_1 be the number of independent cycles in Γ . Then due to the reasons above we have already found a

$$(b_0 - 1) + (n - 1)b_1$$

dimensional family of simple normal forms. What is true, however, is that by this we have already recovered all simple normal forms for f .

5.8. From the discussion above we see that for each $f \in \mathcal{H}_{\Gamma}$ we can define number $S(f) \in \mathbb{N}$ which is the smallest natural number n such that f allows a (simple) (n) -normal form. It is immediate that S gives an additive filtration of $\mathcal{H}_{\Gamma}$, in particular

$$S(f + g) \leq \max(S(f), S(g))$$

with equality if $S(f) \neq S(g)$. What is not so obvious is that it also gives a multiplicative filtration in the sense that

$$S(fg) \leq S(f) + S(g).$$

This, however, follows from Lemma 4.11.

C. Extension to $\overline{\mathbb{C}}$.

5.9. Suppose that $h : \overline{\mathbb{C}} \rightarrow \overline{\mathbb{C}}$ is a conformal map, $\Gamma \subset \mathbb{C}$ is a Lipschitz curve system, and $h(\Gamma) \subset \mathbb{C}$. Suppose that the function $f(z) \in \mathcal{H}_\Gamma$ has an (n) -normal form as above. Then Lemma 4.12.b can be applied and we find that

$$f(z) = \sum_{k=0}^n \int \frac{\tilde{\mu}_k(w)}{(h(w) - h(z))^k}$$

with appropriate measures $\tilde{\mu}_k$. Substituting $z = h^{-1}(s)$ we can immediately see that

$$f(h^{-1}(s)) = \sum_{k=0}^n \int \frac{\tilde{\mu}_k(w)}{(h(w) - s)^k} = \sum_{k=0}^n \int_{\tilde{w} \in h(\Gamma)} \frac{h_* \tilde{\mu}_k(\tilde{w})}{(\tilde{w} - s)^k}.$$

In other terms, the new coefficient measures belong to an (n) -normal form of the function $f \circ h^{-1}$ with respect to the curve system $h(\Gamma)$.

5.10. The argument above, of course, works in a reversed way: if we know that $f \circ h^{-1}$ allows a (simple) (n) -normal form with respect to $h(\Gamma)$ then we may consider a generalized (simple) (n) -form of f with respect to Γ :

Assume that $\Gamma \subset \overline{\mathbb{C}}$ is Lipschitz curve system and $f \in \mathcal{H}_\Gamma$. Then we can define appropriate normal forms as follows: Suppose that $h : \overline{\mathbb{C}} \rightarrow \overline{\mathbb{C}}$ is a conformal map such that $h(\Gamma) \subset \mathbb{C}$. Then $f \circ h^{-1} \in \mathcal{H}_{h(\Gamma)}$, and it allows, a normal form

$$f \circ h^{-1}(z) = \sum_{k=0}^n \int_{\tilde{w} \in h(\Gamma)} \frac{\tilde{\mu}_k(\tilde{w})}{(\tilde{w} - z)^k}.$$

Then by defining the measure $\mu_k = (h^{-1})_* \tilde{\mu}_k$ we find that

$$f(z) = \sum_{k=0}^n \int \frac{\mu_k(w)}{(h(w) - h(z))^k}.$$

So, we see that generalized (simple) normal forms can be defined similarly to (simple) normal forms but $\frac{1}{w-z}$ substituted by $\frac{1}{h(w)-h(z)}$.

5.11. The importance of point 5.9 beside providing inspiration is that it immediately shows that $S(f \circ h^{-1})$ above does not depend on h as long as $h(\Gamma) \subset \mathbb{C}$. Consequently, the filtration S extends to general $\mathcal{H}_\Gamma$, and $S(f)$ is conformal invariant in the sense that $S(f) = S(f \circ h^{-1})$ with h arbitrary.

The concrete form of h is not really important here. In order just to indicate what kind of statements we may obtain this way we state

Proposition 5.12. *Suppose that $\gamma : [a, b] \rightarrow \overline{\mathbb{C}}$ is a Lipschitz curve, $\Gamma = \gamma([a, b])$, $f \in \mathcal{H}_\Gamma$. Suppose that $p \in \overline{\mathbb{C}} \setminus \Gamma$, $t_0 \in [a, b]$. Then for $n+1 \geq S(f)$, we claim, there are unique coefficients $\tilde{c}_j$ and a measure μ on $[a, b]$ such that*

$$f(z) = \sum_{j=0}^n \tilde{c}_j S(\gamma(t_0), z)^j + \int \mu(t) H(\gamma(t), z)^{n+1},$$

where

$$S(\gamma(t_0), z) \equiv \begin{cases} \frac{1}{\gamma(t_0) - z} & \text{if } \gamma(t_0) \in \mathbb{C} \\ z & \text{if } \gamma(t_0) = \infty \end{cases}$$

and

$$H(\gamma(t), z) \equiv \begin{cases} \frac{1}{(z-p)^{-1} - (\gamma(t)-p)^{-1}} & \text{if } p \in \mathbb{C} \\ \frac{1}{\gamma(t_0) - z} & \text{if } p = \infty \end{cases}$$

(ie. a generalized simple normal form).

Analogous statements hold for general Lipschitz curve systems Γ .

Proof. Indeed, the case $p = \infty$ is what we have studied before. The case $p \in \mathbb{C}$ follows from taking $h(z) = \frac{1}{p-z}$. $\square$

D. Preparation for the locality argument.

5.13. In what follows we give a recipe for manipulating certain measures. Although it involves choices the method itself is quite well-defined:

Suppose that $\Gamma \subset \mathbb{C}$ is a Lipschitz curve system, $\overline{\mathbb{C}} \setminus \Gamma = \mathcal{V}_1 \dot{\cup} \mathcal{V}_2$, where $\mathcal{V}_1, \mathcal{V}_2$ are open, and $\mathcal{V}_2$ contains the infinite component of $\overline{\mathbb{C}} \setminus \Gamma$. Assume that $f \in \mathcal{H}_\Gamma$ such that $f|_{\mathcal{V}_2} = 0$. Now suppose that $\mu_0, \dots, \mu_n$ are measures on $\mathbb{C}$ such that they are supported on Γ and

$$f(z) = \sum_{k=0}^n \frac{\mu_k(z)}{(w-z)^k}.$$

Suppose that the connected components of $\mathcal{U} = \text{int } \overline{\mathcal{V}_1}$ are $\mathcal{U}_1, \dots, \mathcal{U}_p$. Our first step among the manipulations is that we cut up the $\mathcal{U}_j$ with some Lipschitz curves such that it yields simply connected Lipschitz domains $\tilde{\mathcal{U}}_j$. We can choose the cutting curves such that they are pairwise disjoint simple curves connecting regular points of Γ . We do that for all j such that $1 \leq j \leq p$. Let $\Gamma' = \Gamma_{\text{ext}} \cup \Gamma_{\text{cut}}$, where $\Gamma_{\text{ext}} = \partial \mathcal{V}_2$ and Γ_{cut} is the supporting set of the cuts. The curve system Γ' has nodes of two type: there are the old nodes coming from Γ and there are the new nodes coming from the cuts.

In the second step, we push the measures $\mu_j|_{\tilde{\mathcal{U}}_j}$ to the boundary and we sum them up. Let

$$g(z) = \int \mu_0 + \sum_{k=1}^n \int \frac{\mu_k|_{\Gamma'}(w)}{(w-z)^k} + \sum_{j=1}^p \sum_{k=1}^n \frac{\tilde{c}_{j,k}}{(\tilde{w}_j - z)^k} + \sum_{k=1}^{n+1} \int \frac{\theta_k(w)}{(w-z)^k},$$

where the latter two terms are obtained according to Lemma 4.15.b. At this point, the supporting set of the measures is Γ' . We know that $f(z) = 0 = g(z)$ for $z \in \mathcal{V}_2$.

In the third step, using appropriate additive reduction we have

$$g(z) = \sum_h \sum_{j=0}^n \frac{c_{j,h}}{(w_h - z)^j} + \int \frac{\mu(w)}{(w-z)^{n+1}},$$

where the w_h are from different connected components of Γ' . Due to residue behaviour the $c_{j,h}$ vanish, so

$$(5) \quad g(z) = \int \frac{\mu(w)}{(w-z)^{n+1}}.$$

Because of the nature of additive reduction and the properties of θ_n we know that locally $\mu(w) = p(w)d\Gamma'(w)$, where $d\Gamma'(w)$ is the variation measure of Γ' with piecewise constant orientation, and $p(w)$ is a piecewise continuous function. We can assume that the discontinuities happen only at the nodes of Γ' .

Lemma 5.14. *Consider*

$$h(z) = \int \frac{\mu(w)}{w - z},$$

which is $n!$ times an n -th primitive function of $g(z)$.

We claim that h extends to the analytic boundary of $\overline{\mathbb{C}} \setminus \Gamma'$. If the $\hat{\gamma}_j$ is the oriented boundary curve of $\tilde{\mathcal{U}}_j$, $\hat{h}_j$ are the corresponding extended values, and γ_j are the underlying curves of $\hat{\gamma}_j$ then

$$(6) \quad g(z) = \sum_{j=1}^p \frac{1}{2\pi i} \int \hat{h}_j(t) \frac{d\gamma_j(t)}{(\gamma(t) - z)^{n+1}}.$$

Proof. First, we claim that $h(z)$ extends continuously to the analytic boundary of $\mathbb{C} \setminus \Gamma'$. Indeed, if s is a regular point of Γ' , then s bounds $\mathcal{V}_2$, on which side $h(z)$ is a polynomial, so by Proposition 3.14 $h(z)$ extends to the other side continuously, too. If s is an old node of Γ' then the argument is similar, except we have to use Proposition 3.17. At new nodes, we claim the behaviour is not far from regular points. Indeed, let us slightly relocate the cut, then we obtain $\tilde{\Gamma}'$ and $\tilde{h}$. Then s is a regular point of $\tilde{\Gamma}'$ and the difference $\tilde{h} - h$ is locally polynomial. This indeed proves the existence of the continuous extension.

Now consider

$$\tilde{h}(z) = \sum_{j=1}^p \frac{1}{2\pi i} \int \hat{h}_j(t) \frac{d\gamma_j(t)}{\gamma(t) - z}.$$

This is equal to $h(z)$ but on $\mathcal{V}_2$ where it vanishes. Nevertheless $\tilde{h} - h$ is a polynomial of degree $n - 1$ on the components of $\mathcal{V}_2$. In particular, taking n -th derivatives and dividing by $n!$ we obtain (6). $\square$

The fourth step of our manipulation process is to replace the normal form (5) by the normal form (6). Notice that this replacement can be done by an additive reduction process, no matter how we argued for the existence of the normal form (6).

6. ARITHMETRICS OF MULTIVALUED LINEAR OPERATORS

6.1. A multivalued linear operator A on a vector space $\mathfrak{V}$ is simply a linear subspace of $\mathfrak{V} \times \mathfrak{V}$. If A is a multivalued linear operator and $a, b, c, d \in \mathbb{C}$, then we may form the multivalued linear operator

$$\frac{aA + b}{cA + d} = \{(ay + bx, cy + dx) : (y, x) \in A\}.$$

This yields an associative $P\mathbb{M}_2(\mathbb{C})$ -action on multivalued linear operators. In simple cases we still use the colloquial notation aA , $A + b$, $b + A$, A^{-1} , $(cA + d)^{-1}$, etc. We have to keep in mind that the definition is a matrix action, so $a + \frac{b}{cA + d}$ is to be imagined as at least three consecutive matrix actions ($A \mapsto A^{-1}$, $B \mapsto \frac{b}{c - B}$, $C \mapsto a + C$) executed but which, naturally, also simplifies according to the associativity of the action. The only possible source of confusion is when $a = c = 0$ because then we have to keep in mind that we started with A . So, in that case it is better to write $\frac{0 \cdot A + b}{0 \cdot A + d}$. If $ad - bc = 0$ then we have a restriction of one of $\alpha \text{Id}_{\mathfrak{V}} = \{(\alpha x, x) : x \in \mathfrak{V}\}$, $\alpha \in \mathbb{C}$ or $\infty \text{Id}_{\mathfrak{V}} = \{(x, 0) : x \in \mathfrak{V}\}$. If $ad - bc \neq 0$ then we have bijective correspondence of multivalued linear operators. If $h(z) = \frac{az + b}{cz + d}$ is the corresponding conformal map on the Riemann sphere, then we use the short notation $h_f(A) = \frac{aA + b}{cA + d}$.

We say that the multivalued linear operator A is an (not necessarily everywhere defined) operator, if $(y, x) \in A, x = 0$ implies $y = 0$, ie. its values are well-defined. In that case we define Ax as y where $(y, x) \in A$.

Lemma 6.2. *a.) If $a, b \in \mathbb{C}, a \neq 0$ then Au makes sense if and only if $(aA + b)u$ makes sense, and in that case $(aA + b)u = aAu + bu$.*

b.) Suppose that Au, Bu make sense, $a, b \in \mathbb{C}$. Then BAu makes sense if and only if $(B + b)(A + a)u$ makes sense, and in that case $(B + b)(A + a)u = abu + bAu + aBu + BAu$.

c.) If Bu and $B^{-1}v$ make sense then $B^{-1}Bu$ and $BB^{-1}v$ make sense, moreover $B^{-1}Bu = u, BB^{-1}v = v$.

d. $(B + b)(B^{-1} + a)u$ makes sense, $a, b \in \mathbb{C}, a \neq 0$ then Bu makes sense. $\square$

Lemma 6.3. *Suppose that $a_1d_1 - b_1c_1 \neq 0, a_2d_2 - b_2c_2 \neq 0, \frac{a_1z+b_1}{c_1z+d_1} \frac{a_2z+b_2}{c_2z+d_2} = \frac{a_3z+b_3}{c_3z+d_3}$. If*

i.) $\frac{a_1A+b_1}{c_1A+d_1}u$ and $\frac{a_2A+b_2}{c_2A+d_2}u$ make sense, or

ii.) $\frac{a_1A+b_1}{c_1A+d_1} \frac{a_2A+b_2}{c_2A+d_2}u$ makes sense, $a_3d_3 - b_3c_3 \neq 0$,

then

$$\frac{a_1A + b_1}{c_1A + d_1} \frac{a_2A + b_2}{c_2A + d_2} u = \frac{a_3A + b_3}{c_3A + d_3} u = \frac{a_2A + b_2}{c_2A + d_2} \frac{a_1A + b_1}{c_1A + d_1} u,$$

including the statement that every expression in the equation makes sense.

Proof. Set $B = \frac{a_1A+b_1}{c_1A+d_1}$. Then $\frac{a_2A+b_2}{c_2A+d_2} = pB^{-1} + q$ with $p, q \in \mathbb{C}, p \neq 0$. What we have to prove is that

$$B(pB^{-1} + q)u = (p + qB)u = (pB^{-1} + q)Bu,$$

including that every term is well-defined.

i.) We know that Bu and $(pB^{-1} + q)u$ exist. In particular, $B^{-1}u$ also exists. Using Lemma 6.2.c we see that $BB^{-1}u$ and $B^{-1}Bu$ also exist, and they are equal to u . Then Lemma 6.2.b implies that the equality above holds.

ii.) In that case $q \neq 0$. Then, according to Lemma 6.2.d Bu exists. By this we have reduced the problem to case (i). $\square$

Lemma 6.4. *Suppose that $w_1, w_2 \in \mathbb{C}, w_1 \neq w_2$. If*

i.) $(w_1 - A)^{-1}u$ and $(w_2 - A)^{-1}u$ make sense, or

ii.) $(w_1 - A)^{-1}(w_2 - A)^{-1}u$ makes sense

then the resolvent identity holds:

$$\begin{aligned} (w_1 - A)^{-1}(w_2 - A)^{-1}u &= (w_2 - w_1)^{-1}((w_1 - A)^{-1}u - (w_2 - A)^{-1}u) \\ &= (w_2 - A)^{-1}(w_1 - A)^{-1}u, \end{aligned}$$

including the statement that all expressions in the identity are well-defined.

In short terms: The resolvent identity holds if either side makes sense.

Proof. Let $B = \frac{w_1 - A}{w_2 - A}, \xi = \frac{1}{w_2 - w_1}$. Then

$$(w_1 - A)^{-1} = \xi(B^{-1} - 1), \quad \text{and} \quad (w_2 - A)^{-1} = \xi(1 - B).$$

In particular, Bu and $B^{-1}u$ exist, hence $B^{-1}Bu$ and $BB^{-1}u$ exist and they are equal to u . We have to prove that

$$\xi(B^{-1} - 1)\xi(1 - B)u = \xi(\xi(B^{-1} - 1)u - \xi(1 - B)u) = \xi(1 - B)\xi(B^{-1} - 1)u$$

ie. that

$$(B^{-1} - 1)(1 - B)u = Bu - 2u + B^{-1}u = (1 - B)(B^{-1} - \text{Id})u.$$

i.) The equality follows from Lemma 6.2.b.

ii.) In that case $q \neq 0$ and the statement is reduced to part (i) by Lemma 6.2.d. $\square$

Ie. the resolvent identity is still valid, which is of crucial importance. It makes possible to generalize the identities concerning rational functions. If $n \geq 0$, $c \in \mathbb{C}$, A is a multivalued linear operator on $\mathfrak{V}$, $u \in \mathfrak{V}$ then we use the notation

$$\frac{c}{A^n}u = c \overbrace{\frac{1}{A} \cdots \frac{1}{A}}^{n \text{ times}} u$$

ie. $\frac{c}{A^n}u$ is A^{-1} iterated n times on u multiplied by the constant c . Now, using the resolvent identity above one can easily prove the full analogue of Lemma 4.1:

Lemma 6.5. *Suppose that $w_1, w_2 \in \mathbb{C}$, $w_1 \neq w_2$. Then*

b.) For $n \in \mathbb{N}$

$$\frac{1}{w_1 - w_2} \left(\frac{1}{(w_1 - A)^n} u - \frac{1}{(w_2 - A)^n} u \right) = - \sum_{j=1}^n \frac{1}{(w_1 - A)^j} \frac{1}{(w_2 - A)^{n-j+1}} u$$

if either side makes sense.

c.) For $n_1, n_2 \in \mathbb{N}$

$$\begin{aligned} \frac{1}{(w_1 - A)^{n_1+1}} \frac{1}{(w_2 - A)^{n_2+1}} u &= \sum_{j=0}^{n_1} \frac{(j + n_2)!}{j! n_2!} \frac{1}{(w_2 - w_1)^{n_2+j+1}} \frac{1}{(w_1 - A)^{n_1-j+1}} u + \\ &+ \sum_{j=0}^{n_2} \frac{(j + n_1)!}{j! n_1!} \frac{1}{(w_1 - w_2)^{n_1+j+1}} \frac{1}{(w_2 - A)^{n_2-j+1}} u \end{aligned}$$

if either side makes sense. □

6.6. Assume from now on that $\mathfrak{V}$ is a locally convex vector space. Suppose that $\Gamma \subset \overline{\mathbb{C}}$ and A is a multivalued linear operator. For $n \in \mathbb{N}$ we define $D_\Gamma^n(A) \subset \mathfrak{V}$ as the set of elements u such that for all $0 \leq m \leq n$ the functions

$$R_\infty : (z_1, \dots, z_m) \in (\Gamma \setminus \{\infty\})^m \mapsto (z_1 - A)^{-1} \cdots (z_m - A)^{-1} u$$

and

$$R_0 : (z_1, \dots, z_m) \in (\Gamma \setminus \{0\})^m \mapsto (z_1^{-1} - A^{-1})^{-1} \cdots (z_m^{-1} - A^{-1})^{-1} u$$

are well-defined and continuous. Notice that $D_{\{\infty\}}^1(A)$ is simply the domain of A if it is an operator. In order to make this notion more friendly, we give another characterization:

Lemma 6.7. *a.) If $\infty \notin \Gamma$ then $u \in D_\Gamma^n(A)$ if and only if for all $0 \leq m \leq n$ the functions*

$$R_\infty : (z_1, \dots, z_m) \in \Gamma^m \mapsto (z_1 - A)^{-1} \cdots (z_m - A)^{-1} u$$

are well-defined and continuous.

b.) If $p \in \mathbb{C}$, $p \notin \Gamma$ then $u \in D_\Gamma^n(A)$ if and only if for all $0 \leq m \leq n$ the functions

$$\begin{aligned} R_p : (z_1, \dots, z_m) \in \Gamma^m \mapsto \\ ((z_1 - p)^{-1} - (A - p)^{-1})^{-1} \cdots ((z_m - p)^{-1} - (A - p)^{-1})^{-1} u \end{aligned}$$

are well-defined and continuous. □

One might think that $u \in D_\Gamma^n(A) \subset \mathfrak{V}$ is characterized by a global property with respect to Γ . This is not the case, it is a local property. Using the resolvent identity it is not hard to prove

Lemma 6.8. *We have $u \in D_\Gamma^n(A)$ if and only if for any $w \in \Gamma$ there is an open set $w \in \mathcal{U}$ such that $u \in D_{\Gamma \cap \mathcal{U}}^n(A)$. $\square$*

Now we can state the generalization of Lemma 4.2. For the sake of simplicity we write $\frac{\theta(t)}{(\gamma(t)-A)^n}u$ instead of $\theta(t)\frac{1}{(\gamma(t)-A)^n}u$. This expression can also be understood either formally or as a vector measure which is the product of the scalar measure $\theta(t)$ and a continuous vector valued function.

Lemma 6.9. *Now assume that $\gamma : [a, b] \rightarrow \mathbb{C}$ is a Lipschitz curve, $\Gamma = \gamma([a, b])$.*

a.) Suppose that $a_1, b_1 \in [a, b]$, $n \geq 1$, $u \in D_\Gamma^{n+1}(A)$. Then

$$\frac{1}{(\gamma(b_1) - A)^n}u - \frac{1}{(\gamma(a_1) - A)^n}u = \int_{t=a_1}^{b_1} \frac{-n d\gamma(t)}{(\gamma(t) - A)^{n+1}}u.$$

b.) Under similar assumption, suppose that

$$\sum_{j=1}^n \frac{c_j}{(-w)^j} \sim \sum_{j=1}^n \frac{c_j^{(\gamma(a_1))}}{(\gamma(a_1) - w)^j} \sim \sum_{j=1}^n \frac{c_j^{(\gamma(b_1))}}{(\gamma(b_1) - w)^j}$$

modulo $O(w^{-n-1})$ as $w \rightarrow \infty$. Then

$$\sum_{j=1}^n \frac{c_j^{(\gamma(b_1))}}{(\gamma(b_1) - A)^j} - \sum_{j=1}^n \frac{c_j^{(\gamma(a_1))}}{(\gamma(a_1) - A)^j} = \int_{t=a_1}^{b_1} \left(\sum_{j=1}^n \binom{n-1}{j-1} c_j \gamma(t)^{n-j} \right) \frac{-n d\gamma(t)}{(\gamma(t) - A)^{n+1}}.$$

c.) Suppose that $a \leq a_1 < b_1 \leq b$, $n_1, n_2 \in \mathbb{N}$, $u \in D_\Gamma^{n_1+n_2+2}(A)$. Then

$$\begin{aligned} & \frac{1}{(\gamma(a_1) - A)^{n_1+1}} \frac{1}{(\gamma(b_1) - A)^{n_2+1}} u = \\ & = \int_{t=a_1}^{b_1} \frac{(n_1 + n_2 + 1)!}{n_1! n_2!} \frac{(\gamma(t) - \gamma(a_1))^{n_2} (\gamma(b_1) - \gamma(t))^{n_1}}{(\gamma(b_1) - \gamma(a_1))^{n_1+n_2+1}} \frac{d\gamma(t)}{(\gamma(t) - A)^{n_1+n_2+2}} u. \end{aligned}$$

Proof. Part (a) follows from Lemma 6.5.a. Part (b) and (c) follow from part (a). $\square$

When we consider superpositions as in Lemma 4.5 and 4.9 it is essential that the integrals in question exist. For this reason we assume that the locally convex vector space $\mathfrak{V}$ is sequentially complete. Proceeding with the generalization of Lemma 4.9:

Theorem 6.10 (Additive reduction). *Suppose that $\Gamma \subset \mathbb{C}$ is a connected Lipschitz curve system, $w_0 \in \Gamma$, $n \in \mathbb{N}$, and the map $s \mapsto \Omega_{w_0, s}^{(\phi)}$ is constructed according to a suitable choice function ϕ . Suppose that μ is a measure on Γ , $u \in D^{n+1}(A)$, $\mathfrak{V}$ is a sequentially closed locally convex vector space. Then*

$$\int \frac{\mu(w)}{(w - A)^n} u = \frac{c_n}{(w_0 - A)^n} u + \int \frac{n \theta(w)}{(w - A)^{n+1}} u,$$

where

$$c_n = \int \mu(s) \quad \text{and} \quad \theta(w) = \int_s \Omega_{w_0, s}^{(\phi)}(w) \mu(s).$$

Proof. It follows from the Fubini integral formulae (4). $\square$

When it comes to multiplication as in Lemma 4.7 and 4.11, in order to generalize them we have to apply for the sequential closedness of A . The reason is that closedness allows to interchange integration and operator actions:

Lemma 6.11. *Suppose that μ is a measure on S , v is a continuous $\mathfrak{V}$ -valued function on S , and A is an operator on $\mathfrak{V}$ such that Av is also continuous on S . Assume that $\mathfrak{V}$ is sequentially complete, A is sequentially closed. Then*

$$A \left(\int v(s) \mu(s) \right) = \int Av(s) \mu(s),$$

including the statement that both sides make sense.

It is a trivial observation that if $a, b, c, d \in \mathbb{C}$, $ad - bc \neq 0$ and A is sequentially closed then $\frac{aA+b}{cA+d}$ is sequentially closed, too. Using that we can state

Theorem 6.12 (Multiplicative reduction). *Suppose that $\Gamma \subset \mathbb{C}$ is a Lipschitz curve system, μ_1 and μ_2 are measures on it. Suppose that the map $(s_1, s_2) \mapsto \Xi_{s_1, s_2; k}^{(\phi, n_1, n_2)}(s)$ is obtained according to a choice function ϕ as earlier. Assume that $u \in D^{n_1+n_2+2}(A)$, $\mathfrak{V}$ is a sequentially closed locally convex vector space, A is a sequentially closed multivalued linear operator. Then*

$$\int \frac{\mu_1(s_1)}{(s_1 - A)^{n_1+1}} \int \frac{\mu_2(s_2)}{(s_2 - A)^{n_2+1}} u = \sum_{k=1}^{n_1+n_2+2} \int \frac{\theta_k(s)}{(s - A)^k} u,$$

where

$$\theta_k(s) = \int_{(s_1, s_2) \in \Gamma \times \Gamma} \Xi_{s_1, s_2; k}^{(\phi, n_1, n_2)}(s) \mu_1(s_1) \mu_2(s_2).$$

Proof. It follows from the previous lemma and the Fubini integral formulae (4). $\square$

The generalization of the rest of Section 4 is straightforward.

Theorem 6.13. *Let $h : \overline{\mathbb{C}} \rightarrow \overline{\mathbb{C}}$ be a conformal map, $h(z) = \frac{az+b}{cz+d}$, $ad - bc \neq 0$.*

a.) *Suppose that $w, h(w) \in \mathbb{C}$. Then*

$$\frac{1}{w - A} = \frac{c}{cw + d} + \frac{ad - bc}{(cw + d)^2} \frac{1}{h(w) - h_f(A)}.$$

b.) *Suppose that $\Gamma \subset \mathbb{C}$ is a Lipschitz curve system, μ is a measure on it. Assume that $h(\Gamma) \subset \mathbb{C}$. Suppose that $u \in D_\Gamma^n(A)$. Then*

$$\int \frac{\mu(w)}{(w - A)^n} u = \sum_{j=0}^n \int \binom{n}{j} \left(\frac{c}{cw + d} \right)^{n-j} \left(\frac{ad - cd}{(cw + d)^2} \right)^j \frac{\mu(w)}{(h(w) - h_f(A))^j} u.$$

$\square$

For the purpose of the following statements let $\mathcal{U} \subset \mathbb{C}$ be a simply connected Lipschitz domain with oriented boundary curve $\hat{\gamma}$ with underlying curve $\gamma : [a, b]_{a \sim b} \rightarrow \mathbb{C}$.

Theorem 6.14. *For any polynomial P with degree $\leq n - 1$, $u \in D_\Gamma^{n+1}(A)$*

$$\int P(\gamma(t)) \frac{d\gamma(t)}{(\gamma(t) - A)^{n+1}} u = 0.$$

Proof. Cutting γ to pieces, it follows from Lemma 6.9.b. $\square$

Theorem 6.15. *Assume that $f : \mathcal{U} \rightarrow \mathbb{C}$ is a holomorphic function which extends to the analytic boundary of $\mathcal{U}$ continuously. Suppose that $\hat{f} : [a, b]_{a \sim b} \rightarrow \mathbb{C}$ gives the*

boundary values in the sense that at $\hat{\gamma}(t)$ the extended value is $\hat{f}(t)$. Suppose that $n \in \mathbb{N}$, $u \in D_{\mathcal{U}}^{n+1}(A)$. Then

$$\int \hat{f}(t) \frac{d\gamma(t)}{(\gamma(t) - A)^{n+1}} u = 0.$$

Proof. If $\tilde{\gamma}$ is a similar curve inside $\mathcal{U}$ then the statement is a consequence of Green's theorem, or rather the fundamental integral formula behind it. Due to the fact that $\mathcal{U}$ is a Lipschitz domain, we can well approximate γ by such $\tilde{\gamma}$, and by continuity the statement follows. $\square$

Theorem 6.16. a.) If $n \geq 1$, $w \in \mathcal{U}$, $t_0 \in [a, b]/\{a \sim b\}$, $u \in D_{\mathcal{U}}^{n+1}(A)$ then

$$\frac{1}{(w - A)^n} u = \frac{1}{(\gamma(t_0) - A)^n} u + n \frac{1}{2\pi i} \int \log(\gamma(t) - w) \frac{d\gamma(t)}{(\gamma(t) - A)^{n+1}} u,$$

where $\log(\gamma(t) - w)$ is a branch such that it is continuous on $([a, b]/\{a \sim b\}) \setminus \{t_0\}$.

b.) If μ is a measure on $\mathcal{U}$, $n \geq 1$, $w \in \mathcal{U}$, $t_0 \in [a, b]/\{a \sim b\}$, $u \in D_{\mathcal{U}}^{n+1}(A)$, then

$$\int \frac{\mu(w)}{(w - A)^n} u = \frac{1}{(\gamma(t_0) - A)^n} \int \mu(w) u + n \frac{1}{2\pi i} \int_{(t,w) \in [a,b] \times \mathcal{U}} r(t, w) \frac{\mu(w) d\gamma(t)}{(\gamma(t) - A)^{n+1}} u$$

as long as i.) for a fixed w the $r(t, w) = \log(\gamma(t) - w)$ is a branch such that it is continuous except at t_0 , ii.) $r(t, w) \mu(w) d\gamma_i(t)$ is Lebesgue integrable.

Proof. a.) By the the standard encircling argument and partial integration

$$(\text{LHS}) = \frac{1}{2\pi i} \int \frac{1}{(\gamma(t) - w)} \frac{d\gamma(t)}{(\gamma(t) - A)^n} = (\text{RHS}).$$

b.) This is just the superposition statement of part (a). $\square$

7. THE HOLOMORPHIC FUNCTIONAL CALCULUS

7.1. Let us fix a Lipschitz curve system $\Gamma \subset \overline{\mathbb{C}}$, a sequentially complete locally convex space $\mathfrak{V}$, and a multivalued linear operator A on $\mathfrak{V}$.

Assume now that $f \in \mathcal{H}_{\Gamma}^n$. Then we define the operator $f(A)$ on $D_{\Gamma}^n(A)$ as follows: Consider any generalized normal form

$$f(z) = \sum_{j=0}^n \int_{w \in \Gamma} \frac{\mu_j(w)}{(h(w) - h(z))^j},$$

where $h : \overline{\mathbb{C}} \rightarrow \overline{\mathbb{C}}$ is a conformal map such that $h(\Gamma) \subset \mathbb{C}$. Then, for $u \in D_{\Gamma}^n(A)$ we define

$$f(A)u = \sum_{j=0}^n \int_{w \in \Gamma} \frac{\mu_j(w)}{(h(w) - h_f(A))^j} u,$$

ie. the evaluation of the normal form for A on u . We have to show that $f(A)u$ above does not depend on the particular choice of the generalized normal form.

Theorem 7.2 (Well-definedness). If $f \in \mathcal{H}_{\Gamma}^n$, $u \in D_{\Gamma}^n(A)$ then

- a.) $f(A)u$ is well-defined.
- b.) $f(A)u$ is linear in f and u .
- c.) If h is a conformal map, then

$$f(A)u = (f \circ h^{-1})(h_f(A))u.$$

d.) If A is sequentially closed then $f(A)$ defines linear maps

$$f(A) : D_{\Gamma}^{n+n_1}(A) \rightarrow D_{\Gamma}^{n_1}(A).$$

Proof. Part (a): First, we consider the case when $h(z) = z$ can be taken. Then we have to show that $f(A)u$ does not depend on the particular choice of the normal form belonging to f . We know that normal forms can be reduced to simple normal forms with fixed poles using Lemma 4.9, while according to Theorem 6.10 the evaluation for A does not change. Then the simple normal forms differ only by some polynomial cycles whose evaluation vanishes according to Theorem 6.14. In the second step, one must show that using different h the evaluation does not change. Indeed, Theorem 6.13.b shows that while obtaining a generalized normal form according to point 5.9 the evaluation of $f(A)u$ does not change. The general case then follows from a formal change of variable $A \mapsto h_f(A)$. Parts (b) and (c) follow from the definition immediately; (d) follows from Lemma 6.11. $\square$

Next we prove the multiplicative property, from which one can develop the spectral mapping theorem, etc.

Theorem 7.3 (Multiplicativity). *Suppose that A is a sequentially closed multivalued linear operator on the sequentially complete locally convex space $\mathfrak{V}$, and $f_1 \in \mathcal{H}_{\Gamma}^{n_1}$, $f_2 \in \mathcal{H}_{\Gamma}^{n_2}$. Then $f_1 f_2 \in \mathcal{H}_{\Gamma}^{n_1+n_2}$, and for $u \in D_{\Gamma}^{n_1+n_2}(A)$*

$$f(A)u = f_1(A)f_2(A)u.$$

Proof. After a conformal change of variable we can assume that $\Gamma \subset \mathbb{C}$. Lemma 4.11 and Theorem 6.12 show that a certain normal form of the product function $f_1(z)f_2(z)$ gives the evaluation $f_1(A)f_2(A)u$. $\square$

Theorem 7.4 (Locality). *Suppose that Γ is a Lipschitz curve system, $\overline{\mathbb{C}} \setminus \Gamma = \mathcal{V}_1 \cup \mathcal{V}_2$, $f \in \mathcal{H}_{\Gamma}^n$ vanishes on $\mathcal{V}_2$, and $u \in D_{\Gamma \cup \mathcal{V}_1}^{n+1}(A)$. Then*

$$f(A)u = 0.$$

Proof. After a conformal change of variable we can assume that $\Gamma \subset \mathbb{C}$. Let us follow the evaluation $f(A)u$ during the preparation argument of 5.13. According to Theorem 6.10 and Theorem 6.16.b we see that the evaluation does not change. Using functional theoretic arguments, in Lemma 5.14, we saw that the prepared normal form can be assumed to be of the very special form (6). But then, according to Theorem 6.15, the evaluation $f(A)u$ vanishes. $\square$

The case of the classical Riesz functional calculus (in case of Lipschitz curves) can be considered as the special case when Γ is a closed simple curve bounding the bounded open set $\mathcal{U}$ which contains the spectrum of A , and $f = F|_{\mathcal{U}}$ where F is holomorphic in a neighborhood of $\overline{\mathcal{U}}$. The locality property is the counterpart of the statement that one can relocate the integrating paths.

8. LOGARITHMS AND FRACTIONAL POWERS

8.1. If we want to consider $\log \frac{aA+b}{cA+d}$ then the natural idea is take the function $f(z) = \widetilde{\log} \frac{az+b}{cz+d} \in \mathcal{H}_{\gamma([a_1, b_1])}$ where $\widetilde{\log}$ is an appropriate branch of logarithm and $\gamma : [a_1, b_1] \rightarrow \overline{\mathbb{C}}$ is a Lipschitz curve starting at $\gamma(a_1) = -d/c$ and ending at $\gamma(b_1) = -b/a$. This prescription is unique except that there is an ambiguity $2\pi i k$ in the logarithm function. Regarding the natural choice we may proceed as follow: Suppose that $\gamma : [a_1, b_1] \rightarrow \overline{\mathbb{C}}$

is a Lipschitz curve and $F : [a_1, b_1] \rightarrow \mathbb{C}z + \mathbb{C}$ is a formal linear function such that $F(t) = u(t)z + v(t)$, where $F(a_1) = cz + d$, $F(b_1) = az + b$, $F(t)$ is a scalar multiple of $\gamma(t) - z$ except at $\gamma(t) = \infty$ and it is a scalar multiple of $1 - \gamma(t)^{-1}z$ except at $\gamma(t) = 0$ (in particular, $-\frac{v(t)}{u(t)} = \gamma(t)$). We assume that the scalar multipliers are locally nonvanishing Lipschitz curves in technical sense. Then, we can define in formal sense

$$\log_{(F)} \frac{az + b}{cz + d} = \int_{t=a_1}^{b_1} \frac{dF(t)}{F(t)} = \int_{t=a_1}^{b_1} \frac{u'(t)z + v'(t)}{u(t)z + v(t)} dt.$$

If we substitute $z \in \mathbb{C} \setminus \gamma([a_1, b_1])$ into z then we obtain a concrete representative $\widetilde{\log} \frac{az+b}{cz+d}$. Also, simplifying the right side one can easily extract a normal form. For $z \in \gamma((a_1, b_1))$ we may define $\log_{(F)} - \frac{az+b}{cz+d}$ as the average of $\log_{(F)} \frac{az^++b}{cz^++d}$ and $\log_{(F)} \frac{az^-+b}{cz^-+d}$.

According to the situation above, for $|\operatorname{Re} \alpha| < 1$ we may define powers as

$$\left(\pm \frac{az + b}{cz + d} \right)_{(F)}^\alpha = \exp \left(\alpha \log_{(F)} \pm \frac{az + b}{cz + d} \right)$$

for $z \in \mathbb{C} \setminus \gamma([a_1, b_1])$ with $(+)$ and for $z \in \gamma((a_1, b_1))$ with $(-)$, respectively.

8.2. A very natural case is when $F(t) = \gamma(t) - z$. This yields the function $\log(\gamma, z)$ and

$$\left(\frac{\gamma(b_1) - z}{\gamma(a_1) - z} \right)_{(\gamma)}^\alpha = 1 + \int_{t=a_1}^{b_1} \left(\frac{\sin \alpha \pi}{\pi} \right) \left(\frac{\gamma(b_1) - \gamma(t)}{\gamma(t) - \gamma(a_1)} \right)^\alpha \frac{d\gamma(t)}{\gamma(t) - z} \in \mathcal{H}_{\gamma([a_1, b_1])}$$

(cf. 2.4). The other very natural case is when we consider the straight segment connecting 1 and z : $F(t) = \frac{1+t}{2} - \frac{1-t}{2}z$ for $t \in [-1, 1]$. The associated logarithm function (in the reverse direction) is

$$\log z = - \int_{t=-1}^1 \frac{1-z}{(1+t) + (1-t)z} dt,$$

ie. the classical logarithm function cut along $\overline{\mathbb{R}^-}$. The associated powers are

$$z^\alpha = 1 - \int_{t=-1}^t \frac{\sin \alpha \pi}{\pi} \left(\frac{1+t}{1-t} \right)^\alpha \frac{(1-z)}{(1+z) + (1-z)t} dt \in \mathcal{H}_{\overline{\mathbb{R}^-}}.$$

(One can pass from $F(t) = t - z$ by substituting $\alpha \rightarrow -\alpha$, $z \rightarrow \frac{z+1}{z-1}$, cf. 2.5.)

Lemma 8.3. *a.) If $n \in \mathbb{N}$, $u \in D_{\gamma([a_1, b_1])}^2(A)$, $g_n(z) = \log(\gamma, z)^n$ then $\log(\gamma, A)$ can be iterated on u arbitrarily many times and*

$$(\log(\gamma, A))^n u = g_n(A)u.$$

b.) Let $n \in \mathbb{N}$, and $S = [-n, n]$ or $S = \{\alpha : |\alpha| < n\}$. If $u \in D_{\gamma([a_1, b_1])}^n(A)$, $f_\alpha(z) = \left(\frac{\gamma(b_1) - z}{\gamma(a_1) - z} \right)_{(\gamma)}^\alpha$ then the function

$$\alpha \in S \mapsto f_\alpha(A)u \in \mathfrak{V}$$

is well-defined and continuous. Moreover, if $n \geq 2$ then

$$f_\alpha(A) = \sum_{n \in \mathbb{N}} \frac{\alpha^n}{n!} g_n(u) = \exp(\alpha \log(\gamma, A))u.$$

c.) It yields $\exp(\alpha_1 \log(\gamma, A)) \exp(\alpha_2 \log(\gamma, A))u = \exp((\alpha_1 + \alpha_2) \log(\gamma, A))u$ as long as $\lceil \alpha_1 \rceil + \lceil \alpha_2 \rceil \leq n$. In particular, if $n \geq 2$ then the $\exp(\alpha \log(\gamma, A))$ form a local group on $D_{\gamma([a_1, b_1])}^n(A)$ with infinitesimal generator $\log(\gamma, A)$.

Proof. For $n \geq 1$

$$g_n(A)u \equiv \int_{t=a_1}^{b_1} \frac{1}{2\pi i} ((\log(\gamma, \gamma(t)) + \pi i)^{n+1} - (\log(\gamma, \gamma(t)) - \pi i)^{n+1}) \frac{d\gamma(t)}{\gamma(t) - A} u.$$

Part (a) follows from multiplicativity. Regarding (b): For $|\alpha| < 1$

$$f_\alpha(A) = 1 + \int_{t=a_1}^{b_1} \left(\frac{\sin \alpha \pi}{\pi} \right) \left(\frac{\gamma(b_1) - \gamma(t)}{\gamma(t) - \gamma(a_1)} \right)_\gamma^\alpha \frac{d\gamma(t)}{\gamma(t) - A}.$$

If $|\alpha| < 1$ then the convergence of the normal form measures implies the statement.

Let $h_n(t) = \frac{1}{2\pi i} ((\log(\gamma, \gamma(t)) + \pi i)^{n+1} - (\log(\gamma, \gamma(t)) - \pi i)^{n+1})$. If $\alpha = \alpha_1 + \dots + \alpha_k$, $|\alpha_i| < 1$ (say $\alpha_i = \alpha/n$) then

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{\alpha^n}{n!} (\log(\gamma, A))^n u &= \sum_{n_1, \dots, n_k=0}^{\infty} \frac{\alpha_1^{n_1}}{n_1!} \dots \frac{\alpha_k^{n_k}}{n_k!} (\log(\gamma, A))^{n_1} \dots (\log(\gamma, A))^{n_k} u = \\ &= \int_{t_1=a_1}^{b_1} \dots \int_{t_k=a_1}^{b_1} \frac{\alpha_1^{n_1}}{n_1!} \dots \frac{\alpha_k^{n_k}}{n_k!} h_{n_1}(t) \dots h_{n_k}(t) \frac{d\gamma(t_k)}{\gamma(t_k) - A} \dots \frac{d\gamma(t_1)}{\gamma(t_1) - A} = \\ &= \exp(\alpha_1 \log(\gamma, A)) \dots \exp(\alpha_k \log(\gamma, A)) u = f_{\alpha_1}(A) \dots f_{\alpha_k}(A) u = f_\alpha(A). \end{aligned}$$

Proceeding backwards we can make sure of convergence. In the last step we collapse the terms belonging to the same sum of powers. That proves the statement for $|\alpha| < n$. For $\alpha \in [-n, n]$ we can argue similarly.

Part (c) follows from the power series expansion. $\square$

8.4. One can make a similar statement for

$$\log A = - \int_{t=-1}^1 \frac{1-A}{(1+t) + (1-t)A} dt.$$

Here for, $|\alpha| < 1$

$$(7) \quad A^\alpha = 1 - \int_{t=-1}^1 \frac{\sin \alpha \pi}{\pi} \left(\frac{1+t}{1-t} \right)^\alpha \frac{1-A}{(1+t) + (1-t)A} dt.$$

Assume now that $\operatorname{Re} \alpha \geq 0$. One can notice that the expression above make sense if u is such that for some $\varepsilon > 0$

$$(1+t)^{\alpha+1-\varepsilon} \frac{1-A}{(1+t) + (1-t)A} u$$

is bounded. We denote the set of such elements u by $D_{\mathbb{R}^-}^{\frac{1}{2}, \alpha}(A)$.

Lemma 8.5. *If $u \in D_{\mathbb{R}^-}^{\frac{1}{2}, \alpha}(A)$ then the function*

$$s \in [-1, 1) \mapsto u_s = \left(1 - \int_{t=s}^1 \frac{\sin \alpha \pi}{\pi} \left(\frac{t-s}{1-t} \right)^\alpha \frac{1-A}{(1+t) + (1-t)A} dt \right) u$$

is continuous, and for $s \in (-1, 1)$

$$u_s = \left(\frac{1+s}{2} + \frac{1-s}{2} A \right)^\alpha u.$$

The choice $s = 1$ yields the same expression as on right side of (7). $\square$

Hence, it is entirely reasonable to define $A^\alpha u$ for $u \in D_{\mathbb{R}^-}^{1,\alpha}(A)$ as in (7). What happened is that we used convergence improvement by singling out a particular normal form, but which idea was supported by a continuity property. Using Lemma 8.5 above, one can easily extend the semigroup property, etc. The formula above already applies to positive operators in Hilbert spaces: If A is positive then $|((1+t) + (1-t)A)^{-1}| \leq \frac{1}{1+t}$, which shows that $D(A) \subset D_{\mathbb{R}^-}^{1,\alpha}(A)$ for $0 \leq \operatorname{Re} \alpha < 1$. Now, (7) is a classical formula ready to relate to other ones usual in the theory of fractional powers.

9. CONVERGENCE IMPROVEMENT

There are many ways to improve the holomorphic calculus above.

9.1. First, one can use more general curve systems, e. g.:

a.) If one examines the proofs before, one can see that instead of Lipschitz curve systems it is sufficient to consider piecewise Lipschitz curve systems where at nodes the sets of curve directions regarding the different pieces are disjoint.

b.) When we consider logarithms and exponentials as in the previous section we can take (possibly closed) piecewise Lipschitz curves. For example, Lemma 3.12 shows that logarithmic singularities are rather nice, and by deleting smaller and smaller neighborhoods of the nodes taking limits allow us, by continuity, to pass from disjoint unions of Lipschitz curves to piecewise Lipschitz curves.

There are also many possibilities to relax analytic environment, e. g.:

c.) Instead of allowing all possible normal forms one can restrict to simple normal forms with, say, $\mathcal{L}^1$ measure terms. In this case the condition $u \in D_\Gamma^n(A)$ can be relaxed, principally the iterated resolvent does not need to be defined at the diagonal of Γ^n .

d.) More generally, various growth conditions may help to extend the domains of the operators. For example, it is useful to require only some polynomial growth conditions as in 8.4.

REFERENCES

- [1] C. A. Berenstein, R. Gay: *Complex variables. An introduction*. Graduate Texts in Mathematics 125. Springer-Verlag, New York, Berlin, Heidelberg, 1991.
- [2] N. Dunford, J. T. Schwartz: *Linear Operators, Part I: General Theory*. Interscience Publishers, New York, 1967.
- [3] M. Haase: *The Functional Calculus for Sectorial Operators*. Operator Theory: Advances and Applications 169. Birkhäuser Verlag, Basel, 2006.
- [4] H. Komatsu: Fractional powers of operators. *Pacific J. Math.* **19** (1966), 285–346.
- [5] Gy. Lakos: Notes on Lebesgue integration. arXiv:math.FA/0506185.
- [6] Gy. Lakos: Elementary spectral calculations in rings. arXiv:math.RA/0610921.
- [7] C. Martínez Carracedo, M. Sanz Alix: *The theory of fractional powers of operators*. North-Holland Mathematics Studies, 187. North-Holland, Amsterdam, 2001.

DEPARTMENT OF GEOMETRY, EÖTVÖS UNIVERSITY, PÁZMÁNY PÉTER S. 1/C, BUDAPEST, H-1117, HUNGARY

E-mail address: lakos@cs.elte.hu